

## DECISION MAKING UNDER INCOMPLETE DATA: INTUITIONISTIC MULTI FUZZY IDEALS OF NEAR-RING APPROACH

Nadia Batool<sup>1\*</sup>, Sadaqat Hussain<sup>1</sup>, Nasreen Kausar<sup>2</sup>, Mohammed Munir<sup>3</sup>,  
Rita Yi Man Li<sup>4\*</sup> and Salma Khan<sup>5</sup>

<sup>1</sup>Department of Mathematics, University of Baltistan Skardu, Gilgit Baltistan 16100,  
Pakistan,

<sup>2</sup>Department of Mathematics, Faculty of Arts and Sciences, Yildiz Technical University,  
Esenler 34220, Istanbul, Turkey,

<sup>3</sup>Department of Mathematics, Government Postgraduate College, Abbottabad, Pakistan,

<sup>4</sup>Sustainable Real Estate Research Center, Department of Economics and Finance, Hong  
Kong Shue Yan University, Hong Kong 999077, China

<sup>5</sup>Department of Mathematics and Statistics, Hazara University Mansehra 21120,  
Khyber Pakhtunkhwa, Pakistan.

Received: 10 September 2022;

Accepted: 30 November 2022;

Available online: 4 January 2023.

*Original Scientific Paper*

**Abstract.** *Real-world data is often partial, uncertain, or incomplete. Decision making based on data as such can be addressed by fuzzy sets and related systems. This article studies the intuitionistic multi-fuzzy sub-near rings and Intuitionistic multi-fuzzy ideals of near rings. It presents some of the elementary operations and relations defined on these structures. The concept of level subsets and support of the Intuitionistic multi-fuzzy sub-near ring is also presented. It looks into and demonstrated a few characteristics of intuitionistic multi-fuzzy near-rings and ideals. This research advances fuzzy set theory, which is often applied to problems involving pattern recognition and multiple criterion decision-making. Thus, the results may be beneficial to artificial intelligence related research. Alternatively, the intuitionistic multi-fuzzy approach may be applied to vector spaces and modules or extended to inter-valued fuzzy systems.*

**Keywords:** *Intuitionistic Fuzzy Set, Near-ring, Fuzzy Multi Near-ring, Intuitionistic multi fuzzy Near-ring, Ideals, fuzzy multi ring*

\* Corresponding author.

Email addresses: nadiabatool1512@gmail.com (N. Batool), sadaqatku@gmail.com (S. Hussain), kausar.nasreen57@gmail.com (N. Kausar), dr.mohammadmunir@gmail.com (M. Munir), ritarec1@yahoo.com.hk (Y.M.L.Rita), salmakhan359@gmail.com (S. Khan)

## 1. Introduction

The concept of "fuzzy sets" was initially proposed by Zadeh (Zadeh, 1965) (Zadeh, 1965), which opened the door for set theory researchers. Many versions and generalizations of fuzzy sets have appeared to solve problems such as multi-criteria decision-making, recognition of patterns and diagnosis of diseases (Broumi, Ajay, et al., 2022) (Ashraf et al., 2022; Broumi, Sundareswaran, et al., 2022) waste management (Zhumadillayeva et al., 2020) and voltage balancing (Taghieh et al., 2022). Applications of the fuzzy systems to some other areas can be studied (Gulzar, Alghazzawi, et al., 2020; Gulzar, Mateen, et al., 2020; Kausar, 2019; Kausar et al., 2020; Riaz et al., 2022b). Due to the appearance of fuzzy sets and their associated systems as problem-solving tools in MCDM (Abbas et al., n.d.; Abdullah, 2013; Kahraman, 2008) and other disciplines of our daily life, more people were attracted towards this area of research. Theoretical mathematicians use this idea to generalize well-known mathematical structures. For example, Rosenfeld used it to develop a fuzzy group structure and generalize the classical group. This was the initial development of fuzzy group theory. Liu introduced fuzzy rings and studied various properties of rings and ideals in a fuzzy context. Within a few years, the fuzzification of algebraic structures became a hot topic within the research community. Researchers developed more fuzzy algebraic structures like fuzzy modules, fuzzy algebras, fuzzy sub near rings etc. were developed [see (Al-Husban, 2021; Fathi & Salleh, 2009; Hur et al., 2005; Rahman & Saikia, 2012; Zhan & Ma, 2005)]. Salah Abuzaid was the first who introduced the notion of fuzzy sub near-rings (Abou-Zaid, 1991) and later, its variants were studied (Asif et al., 2020; Hussain et al., 2022).

The concept of multisets is initiated by Yager (Yager, 1986). The multi-fuzzy groups were proposed by T. K. Shinoj et al. [see (Dresher & Ore, 1938; Shinoj et al., 2015)]. They studied the basic properties of multi-fuzzy groups and presented a few preliminary results. The fuzzy versions of multi-subrings and their ideals were established by L. Sujatha (Sujatha, 2014) in 2014. She also proved that the finite multi-fuzzy subrings (ideals) intersection is a multi-fuzzy subring (ideal). Fuzzy multi-near-rings and their associated multi-ideals were introduced by Tahan et al. (Al Tahan et al., 2021) in 2021. They defined various operations on multi-ideals of fuzzy near-rings. They presented foundational results related to fuzzy multi sub-near-rings, fuzzy multi-ideals and the operations defined on multi-ideals. The anti-fuzzy multi-ideals of near-rings were considered by Hoskova (Hoskova-Mayerova & Al Tahan, 2021).

One of the well-known generalizations of fuzzy sets is the intuitionistic fuzzy set proposed by Atanassov (Atanassov, 1986). Renowned mathematicians also use this set to fuzzify algebraic structures. Fathi was the first to describe the notion intuitionistic fuzzy group (Fathi & Salleh, 2009) Consequently, the intuitionistic versions of groups, rings, ideals, modules, near-rings etc., have been established (Hur et al., 2005). Many researchers (Kausar & Waqar, 2019; Kousar et al., 2021, 2022; Riaz et al., 2022a) used intuitionistic fuzzy sets and developed different structures. These included triangular intuitionistic fuzzy linear programming, lattice-valued intuitionistic fuzzy subgroup type-3, algebraic codes over lattice-valued intuitionistic fuzzy type-3 submodules, codes over lattice-valued intuitionistic fuzzy set type-3, non-associative ordered semi-groups by intuitionistic fuzzy bi ideals. Others further developed different versions of intuitionistic fuzzy sets, such as complex intuitionistic fuzzy sets in group theory and t-intuitionistic fuzzy subgroups (Gulzar, Alghazzawi, et al., 2020; Gulzar, Mateen, et al., 2020). It also encompasses the finite

Decision making under incomplete data: Intuitionistic multi fuzzy ideals of near-ring approach intuitionistic anti-fuzzy normal subrings' direct product (Kausar, 2019) intuitionistic fuzzy normal subrings (Shah et al., 2012).

The primary purpose of this study is to find the connections between multi-fuzzy near-rings and intuitionistic fuzzy sets. For this, we first define intuitionistic fuzzy multi-near-rings (IFMNRs) and ideals associated with this structure. We also define the basic operations and study the critical properties of this structure. Moreover, we study the support and  $(\alpha, \beta)$  – level subsets of IFMNRs and produce a few related results. This work contributes to the fuzzy set theory and fuzzy algebra, which are extensively used to solve multi-criteria decision-making and pattern recognition problems (Kaharman et. al., 2008).

## 2. Motivation and Scope

Fuzzy sets and associated systems deal with theoretical and practical problems equipped with incomplete, uncertain or ambiguous data. Although this is a very young discipline started with the work of L. A. Zadeh but gained the attention of many researchers quickly. The reason is that it can be directly applied to theoretical and daily life problems, including Multi-criteria decision-making, Pattern recognition, disease diagnosis and Management. So it is worth studying the fuzzy systems to produce more sufficient and adequate theoretical bases used to develop better problem-solving tools. This article also proposes a fuzzy algebraic system intuitionistic fuzzy multi-near-rings, a generalization of well-known fuzzy multi-near-rings.

## 3. Preliminaries

This study first recalls some basic definitions of fuzzy and multi-fuzzy sets.

**Definition 2.1:** Let  $\mathcal{N}$  be a non-empty set, then a fuzzy set  $\tilde{\mathcal{A}}$  is given by an ordered pair (Zadeh, 1965)

$$\tilde{\mathcal{A}} = \{(s, \Omega_{\tilde{\mathcal{A}}}(s)) / s \in \mathcal{N}\}$$

$\Omega_{\tilde{\mathcal{A}}}$  is the degree of membership, and  $\Omega_{\tilde{\mathcal{A}}} : \mathcal{N} \rightarrow [0,1]$  is the membership function.

**Example 1.** If  $\mathcal{N} = \{s, t, z\}$  then  $\tilde{\mathcal{A}} = \{(s, 0.51), (t, 0.03), (z, 0.21)\}$  be fuzzy set of  $\mathcal{N}$ .

**Definition 2.2:** Let  $\mathcal{N}$  be a set that is not empty. An intuitionistic fuzzy set will be (Atanassov, 1986)

$$\tilde{\mathcal{A}} = \{ \langle s, \Omega_{\tilde{\mathcal{A}}}(s), \mathcal{U}_{\tilde{\mathcal{A}}}(s) \rangle / s \in \mathcal{N} \}$$

Where  $\Omega_{\tilde{\mathcal{A}}}$  and  $\mathcal{U}_{\tilde{\mathcal{A}}}$  are the degrees of membership and non-membership function, respectively, they are defined as  $\Omega_{\tilde{\mathcal{A}}} : \mathcal{N} \rightarrow [0,1]$  and  $\mathcal{U}_{\tilde{\mathcal{A}}} : \mathcal{N} \rightarrow [0,1]$ .

For each  $\epsilon \in \mathcal{N}$   $0 \leq \Omega_{\tilde{\mathcal{A}}}(s) + \mathcal{U}_{\tilde{\mathcal{A}}}(s) \leq 1$ .

**Remark 1.** Every fuzzy set is an intuitionistic fuzzy set.

**Definition 2.3:** Let  $\mathcal{N}$  be a non-empty set, then  $\mathcal{M}$  be a multiset drawn from  $\mathcal{N}$ , characterized by a count function  $\mathcal{CM} : \mathcal{N} \rightarrow \mathbb{N}$ , where  $\mathcal{CM}$  represent the number of repetition of an element in  $\mathcal{M}$  and  $\mathbb{N}$  is set of positive integers (Yager, 1987, Hoskova et. al., 2020).

Let  $\mathcal{N} = \{s_1, s_2, s_3, \dots, s_n\}$  be a set; then a multiset  $\mathcal{M}$  will be represented as  $\mathcal{M} = \{m_1/s_1, m_2/s_2, m_3/s_3, \dots, m_n/s_n\}$  where  $m_i = \{m_{i1}, m_{i2}, m_{i3}, \dots, m_{in}\}$  represents the number of repetition of an element in  $\mathcal{M}$ .

**Example 2.** Let  $\mathcal{N} = \{s, t, z\}$  be a set, then  $\mathcal{M} = \{s, t, t, s, z, t, t\}$  is a multiset, and it can also be represented as  $\mathcal{M} = \{2/s, 4/t, 1/z\}$ .

**Definition 2.4:** Per Tahan et al. (2021), let  $\mathcal{N}$  be a non-empty set then a fuzzy multiset constructed from  $\mathcal{N}$  can be represented as;

$$\mathcal{A} = \{(s, \mathcal{CM}_{\mathcal{A}}(s)) / s \in \mathcal{N}\}$$

Where  $\mathcal{CM}_{\mathcal{A}}: \mathcal{N} \rightarrow Q$  represents the count membership function, here  $Q$  is a set of all crisp multisets which is constructed from unit interval  $[0,1]$  and for each  $s \in \mathcal{N}$ ,  $\mathcal{CM}_{\mathcal{A}}(s)$  is a decreasingly ordered sequence that is  $\Omega_{\mathcal{A}}^1(s) \geq \Omega_{\mathcal{A}}^2(s) \geq \Omega_{\mathcal{A}}^3(s) \geq \dots \geq \Omega_{\mathcal{A}}^n(s)$ .

**Definition 2.5:** Let  $\mathcal{N}$  be a non-empty set. An intuitionistic version of a fuzzy multiset can be represented (Onasanya et. al., 2019);

$$\mathcal{A} = \{< s, \mathcal{CM}_{\mathcal{A}}(s), \mathcal{CN}_{\mathcal{A}}(s) > / s \in \mathcal{N}\}$$

$\mathcal{CM}_{\mathcal{A}}: \mathcal{N} \rightarrow Q$  represents the count membership function, and  $\mathcal{CN}_{\mathcal{A}}: \mathcal{N} \rightarrow Q$  shows the count non-membership function. Here  $Q$  is a set of all crisp multisets which is constructed from unit interval  $[0,1]$ , and for each  $s \in \mathcal{N}$ ,  $\mathcal{CM}_{\mathcal{A}}(s)$  is a decreasingly ordered sequence that is  $\Omega_{\mathcal{A}}^1(s) \geq \Omega_{\mathcal{A}}^2(s) \geq \Omega_{\mathcal{A}}^3(s) \geq \dots \geq \Omega_{\mathcal{A}}^n(s)$  and  $\mathcal{CN}_{\mathcal{A}}(s)$  is denoted by  $(\mathcal{U}_{\mathcal{A}}^1(s), \mathcal{U}_{\mathcal{A}}^2(s), \mathcal{U}_{\mathcal{A}}^3(s), \dots, \mathcal{U}_{\mathcal{A}}^n(s))$ . For each  $s \in \mathcal{N}$ ,  $0 \leq \mathcal{CM}_{\mathcal{A}}(s) + \mathcal{CN}_{\mathcal{A}}(s) \leq 1$ .

**Remark 2.**  $\mathcal{CM}_{\mathcal{A}}(s)$  is ordered decreasingly, but the corresponding  $\mathcal{CN}_{\mathcal{A}}(s)$  may need to be in decreasing and increasing order.

**Remark 3.** An intuitionistic fuzzy set on set  $\mathcal{N}$  can be treated as a particular case of an intuitionistic fuzzy multiset, if  $\mathcal{CM}_{\mathcal{A}}(s) = \Omega_{\mathcal{A}}^1(s)$ .  $\mathcal{CN}_{\mathcal{A}}(s) = \mathcal{U}_{\mathcal{A}}^1(s) \forall s \in \mathcal{N}$ .

**Example 3.** Let  $\mathcal{N} = \{a, b, c\}$ . Then,  $\mathcal{A}$  is an intuitionistic fuzzy multiset over  $\mathcal{N}$  with count functions:

$$\mathcal{CM}_{\mathcal{A}}(s) = \begin{cases} 0.4, 0.4, 0.6 & \text{if } s = a \\ 0.1, 0.1, 0.1 & \text{if } s = b \\ 1, 1, 0.5 & \text{if } s = c \end{cases}$$

$$\mathcal{CN}_{\mathcal{A}}(s) = \begin{cases} 0.6, 0.6, 0.4 & \text{if } s = a \\ 0.9, 0.9, 0.9 & \text{if } s = b \\ 0.5 & \text{if } s = c \end{cases}$$

**Definition 2.6:** According to Tahan et. al (2021), Shinoj et. al., (2015), Hoskova et. al. (2021), let  $\mathcal{N} \neq \emptyset$  be a set,  $\mathcal{A}$  and  $\mathcal{B}$  be two intuitionistic fuzzy multisets over  $\mathcal{N}$  with fuzzy count functions  $\mathcal{CM}_{\mathcal{A}}(s)$ ,  $\mathcal{CM}_{\mathcal{B}}(s)$  and  $\mathcal{CN}_{\mathcal{A}}(s)$ ,  $\mathcal{CN}_{\mathcal{B}}(s)$  respectively, then:

$\mathcal{A} \subseteq \mathcal{B}$  if  $\mathcal{CM}_{\mathcal{A}}(s) \leq \mathcal{CM}_{\mathcal{B}}(s)$  and  $\mathcal{CN}_{\mathcal{A}}(s) \leq \mathcal{CN}_{\mathcal{B}}(s) \forall s \in \mathcal{N}$

$\mathcal{A} = \mathcal{B}$  if  $\mathcal{CM}_{\mathcal{A}}(s) = \mathcal{CM}_{\mathcal{B}}(s)$  and  $\mathcal{CN}_{\mathcal{A}}(s) = \mathcal{CN}_{\mathcal{B}}(s) \forall s \in \mathcal{N}$

$\mathcal{A} \cap \mathcal{B}$  is defined as;

$\mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}(s) = \min \{\mathcal{CM}_{\mathcal{A}}(s), \mathcal{CM}_{\mathcal{B}}(s)\}$  and  $\mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}(s) = \max \{\mathcal{CN}_{\mathcal{A}}(s), \mathcal{CN}_{\mathcal{B}}(s)\}$

$\mathcal{A} \cup \mathcal{B}$  is defined as;

$\mathcal{CM}_{\mathcal{A} \cup \mathcal{B}}(s) = \max \{\mathcal{CM}_{\mathcal{A}}(s), \mathcal{CM}_{\mathcal{B}}(s)\}$  and  $\mathcal{CN}_{\mathcal{A} \cup \mathcal{B}}(s) = \min \{\mathcal{CN}_{\mathcal{A}}(s), \mathcal{CN}_{\mathcal{B}}(s)\}$

The complement of an intuitionistic fuzzy multiset is defined as;

$$\mathcal{A}^c = \{< s, \mathcal{CN}_{\mathcal{A}}(s), \mathcal{CM}_{\mathcal{A}}(s) > / s \in \mathcal{N}\}$$

**Example 4.** Let  $\mathcal{N} = \{a, b, c, d\}$  and  $\mathcal{A}, \mathcal{B}$  be two intuitionistic fuzzy multisets over  $\mathcal{N}$  that is;

Decision making under incomplete data: Intuitionistic multi fuzzy ideals of near-ring approach

$\mathcal{A} = \{ \langle a, (0.8, 0.8, 0.6), (0.2, 0.2, 0.4) \rangle, \langle b, (0.7, 0.7, 0.7), (0.3, 0.3, 0.3) \rangle, \langle c, (1, 0.9), (0.1) \rangle \}$ ,

$\mathcal{B} = \{ \langle a, (1, 0.5, 0.5), (0.5, 0.5) \rangle, \langle c, (0.6, 0.6), (0.4, 0.4) \rangle \}$  Then;

$\mathcal{A} \cap \mathcal{B} = \{ \langle a, (0.8, 0.5, 0.5), (0.2, 0.5, 0.5) \rangle, \langle c, (0.6, 0.6), (0.4, 0.4) \rangle \}$

And;

$\mathcal{A} \cup \mathcal{B} = \{ \langle a, (1, 0.8, 0.6), (0.2, 0.4) \rangle, \langle c, (1, 0.9), (0.1) \rangle \}$ .

**Definition 2.7:** Let  $\mathcal{N}$  be a non-empty set, then  $(\mathcal{N}, +, \cdot)$  is said to be left (right) *NR* if (Tahan et. al., 2021, Asif, et. al., 2020, Hussain et. al., 2022);

$(\mathcal{N}, +)$  is a group;

$(\mathcal{N}, \cdot)$  is a semi-group;

$\mathcal{N}$  Satisfies left (right) distributive law, that is;

$s \cdot (t + z) = s \cdot t + s \cdot z \forall s, t, z \in \mathcal{N}$  (Left distributive)

$(t + z) \cdot s = t \cdot s + z \cdot s \forall s, t, z \in \mathcal{N}$  (Right distributive)

**Example 5.** Let  $\mathbb{R}$  be a set of real numbers, then  $(\mathbb{R}, +, \cdot)$  form *NR* under standard "+" and multiplication is defined as  $a \cdot b = a \forall a, b \in \mathbb{R}$ .

**Note:** Throughout this text, we write *NR* instead of near-ring and *NRs* for near-rings.

**Definition 2.8:** Let  $(\mathcal{N}, +, \cdot)$  be a *NR*, and  $\mathcal{J}$  be a sub-*NR* of  $\mathcal{N}$  then  $\mathcal{J}$  is said to be an ideal of  $\mathcal{N}$  (Tahan et. al., 2021, Asif, et. al., 2020, Hussain et. al., 2022), if;

- i.  $s + t - s \in \mathcal{J} \forall s \in \mathcal{N} t \in \mathcal{J}$
- ii.  $s \cdot t \in \mathcal{J} \forall s \in \mathcal{N} \wedge t \in \mathcal{J}$
- iii.  $(s + a) \cdot t - s \cdot t \in \mathcal{J} \forall s, t \in \mathcal{N} \wedge a \in \mathcal{J}$

**Remark 4** If  $\mathcal{J}$  fulfils conditions (i) and (ii), then  $\mathcal{J}$  is said to be the left ideal, and if  $\mathcal{J}$  satisfies (i) and (iii), then  $\mathcal{J}$  is said to be the ideal of  $\mathcal{N}$ . If  $\mathcal{J}$  is the left and right ideal of  $\mathcal{N}$ , then  $\mathcal{J}$  is said to be an ideal of  $\mathcal{N}$ .

**Example 6.** Let  $\mathcal{M}_2(\mathbb{Z})$  be the set of all possible 2 by 2 matrices with entries from  $\mathbb{Z}$  and  $\mathcal{M}_2(\mathbb{Z})$  form *NR* under standard addition and multiplication is defined as  $(a_{ij}) \cdot (b_{ij}) = (a_{ij})$  for some  $(a_{ij}), (b_{ij}) \in \mathcal{M}_2(\mathbb{Z})$  then  $\mathcal{M}_2(2\mathbb{Z})$  is right ideal of  $\mathcal{M}_2(\mathbb{Z})$ .

**Definition 2.9:** (Abou-Zaid, 1991) Let  $(\mathcal{N}, +, \cdot)$  be a *NR* and  $\mathcal{A}$  be fuzzy set over  $\mathcal{N}$  then  $\mathcal{A}$  is said to be fuzzy sub-*NR* of  $\mathcal{N}$  if  $\forall s, t \in \mathcal{N}$  the following condition satisfies;

- I.  $\Omega_{\mathcal{A}}(s - t) \geq \Omega_{\mathcal{A}}(s) \wedge \Omega_{\mathcal{A}}(t)$
- II.  $\Omega_{\mathcal{A}}(s \cdot t) \geq \Omega_{\mathcal{A}}(s) \wedge \Omega_{\mathcal{A}}(t)$

**Definition 2.10:** (Abou-Zaid, 1991) Let  $(\mathcal{N}, +, \cdot)$  be a *NR* and  $\mathcal{A}$  be fuzzy set over  $\mathcal{N}$  then  $\mathcal{A}$  is said to be the fuzzy ideal of *NR* if  $\forall s, t \in \mathcal{N}$  the following condition satisfies;

- I.  $\Omega_{\mathcal{A}}(s - t) \geq \Omega_{\mathcal{A}}(s) \wedge \Omega_{\mathcal{A}}(t)$
- II.  $\Omega_{\mathcal{A}}(s \cdot t) \geq \Omega_{\mathcal{A}}(s) \wedge \Omega_{\mathcal{A}}(t)$
- III.  $\Omega_{\mathcal{A}}(s + t - s) \geq \Omega_{\mathcal{A}}(t)$
- IV.  $\Omega_{\mathcal{A}}(s \cdot t) \geq \Omega_{\mathcal{A}}(t)$
- V.  $\Omega_{\mathcal{A}}((s + a)t - st) \geq \Omega_{\mathcal{A}}(a) \forall a \in \mathcal{N}$

**Definition 2.11:** (Tahan et. al., 2021) Let  $(\mathcal{N}, +, \cdot)$  be a *NR* and  $\mathcal{A}$  be fuzzy multiset over  $\mathcal{N}$  then  $\mathcal{A}$  is said to be fuzzy multi sub-*NR* of  $\mathcal{N}$  if  $\forall s, t \in \mathcal{N}$  the following condition satisfies;

- I.  $\mathcal{CM}_{\mathcal{A}}(s - t) \geq \mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{A}}(t)$
- II.  $\mathcal{CM}_{\mathcal{A}}(s \cdot t) \geq \mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{A}}(t)$

**Definition 2.12:** (Tahan et. al., 2021) Let  $(\mathcal{N}, +, \cdot)$  be a *NR* and  $\mathcal{A}$  be a fuzzy multiset over  $\mathcal{N}$  then  $\mathcal{A}$  is said to be a fuzzy multi ideal of *NR*  $\mathcal{N}$  if  $\forall s, t \in \mathcal{N}$  the following condition satisfies;

- III.  $\mathcal{CM}_{\mathcal{A}}(s - t) \geq \mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{A}}(t)$
- IV.  $\mathcal{CM}_{\mathcal{A}}(s \cdot t) \geq \mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{A}}(t)$
- V.  $\mathcal{CM}_{\mathcal{A}}(s + t - s) \geq \mathcal{CM}_{\mathcal{A}}(t)$
- VI.  $\mathcal{CM}_{\mathcal{A}}(s \cdot t) \geq \mathcal{CM}_{\mathcal{A}}(t)$
- VII.  $\mathcal{CM}_{\mathcal{A}}((s + a)t - st) \geq \mathcal{CM}_{\mathcal{A}}(a) \forall a \in \mathcal{N}$

#### 4. Main Result

**Definition 3.1:** Let  $(\mathcal{N}, +, \cdot)$  be a *NR*. An intuitionistic multi-fuzzy set  $\mathcal{A}$  is an intuitionistic multi-fuzzy sub-*NR* over  $\mathcal{N}$ . If  $\forall s, t \in \mathcal{N}$ , the following conditions are satisfied;

- I.  $\mathcal{CM}_{\mathcal{A}}(s - t) \geq \mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{A}}(t)$  and  $\mathcal{CN}_{\mathcal{A}}(s - t) \leq \mathcal{CN}_{\mathcal{A}}(s) \vee \mathcal{CN}_{\mathcal{A}}(t)$
- II.  $\mathcal{CM}_{\mathcal{A}}(s \cdot t) \geq \mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{A}}(t)$  and  $\mathcal{CN}_{\mathcal{A}}(s \cdot t) \leq \mathcal{CN}_{\mathcal{A}}(s) \vee \mathcal{CN}_{\mathcal{A}}(t)$

Where  $\mathcal{CM}_{\mathcal{A}}$  and  $\mathcal{CN}_{\mathcal{A}}$  count membership and count non-membership functions, respectively.

**Example 7.** Let  $(\mathbb{Z}, +, \cdot)$  be a *NR* under standard addition and multiplication. Then,  $\mathcal{A}$  is an intuitionistic multi-fuzzy sub-*NR* over  $\mathbb{Z}$  with fuzzy count functions given by;

$$\mathcal{CM}_{\mathcal{A}} = \begin{cases} (0.21, 0.21, 0.21) & \text{if } s \text{ even number} \\ 0 & \text{otherwise} \end{cases}$$

$$\mathcal{CN}_{\mathcal{A}} = \begin{cases} (0.79, 0.79, 0.79) & \text{if } s \text{ even number} \\ 1 & \text{otherwise} \end{cases}$$

**Definition 3.2:** Let  $(\mathcal{N}, +, \cdot)$  be a *NR*. An intuitionistic multi-fuzzy set  $\mathcal{A}$  is considered an intuitionistic fuzzy multi-ideal of  $\mathcal{N}$ . If  $\forall s, t \in \mathcal{N}$ , the following conditions are satisfied;

$$\mathcal{CM}_{\mathcal{A}}(s - t) \geq \mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{A}}(t) \text{ and } \mathcal{CN}_{\mathcal{A}}(s - t) \leq \mathcal{CN}_{\mathcal{A}}(s) \vee \mathcal{CN}_{\mathcal{A}}(t)$$

$$\mathcal{CM}_{\mathcal{A}}(s \cdot t) \geq \mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{A}}(t) \text{ and } \mathcal{CN}_{\mathcal{A}}(s \cdot t) \leq \mathcal{CN}_{\mathcal{A}}(s) \vee \mathcal{CN}_{\mathcal{A}}(t)$$

$$\mathcal{CM}_{\mathcal{A}}(s + t - s) \geq \mathcal{CM}_{\mathcal{A}}(t) \text{ and } \mathcal{CN}_{\mathcal{A}}(s + t - s) \leq \mathcal{CN}_{\mathcal{A}}(t)$$

$$\mathcal{CM}_{\mathcal{A}}(s \cdot t) \geq \mathcal{CM}_{\mathcal{A}}(t) \text{ and } \mathcal{CN}_{\mathcal{A}}(s \cdot t) \leq \mathcal{CN}_{\mathcal{A}}(t)$$

$$\mathcal{CM}_{\mathcal{A}}((s + a)t - st) \geq \mathcal{CM}_{\mathcal{A}}(a) \text{ and}$$

$$\mathcal{CN}_{\mathcal{A}}((s + a)t - st) \leq \mathcal{CN}_{\mathcal{A}}(a) \quad \forall a \in \mathcal{N}$$

**Example 8.** Let  $(\mathbb{Z}, +, \cdot)$  be a *NR* under standard addition and multiplication. Then,  $\mathcal{A}$  is an intuitionistic multi-fuzzy ideal of  $\mathbb{Z}$  with fuzzy count functions given by;

$$\mathcal{CM}_{\mathcal{A}} = \begin{cases} (1, 0.9, 0.7, 0.5, 0.4) & \text{if } s \text{ is a multiple of } 3 \\ (0.9, 0.6, 0.3) & \text{otherwise} \end{cases}$$

$$\mathcal{CN}_{\mathcal{A}} = \begin{cases} (0.1, 0.3, 0.5, 0.6) & \text{if } s \text{ is multiple of } 3 \\ (0.1, 0.4, 0.7) & \text{otherwise} \end{cases}$$

**Remark 5.** Every intuitionistic multi-fuzzy ideal of a *NR*  $\mathcal{N}$  is an intuitionistic multi-fuzzy sub-*NR* of  $\mathcal{N}$ .

This study then demonstrates some exciting results for intuitionistic multi-fuzzy near-rings, which are proven for other algebraic structures (Al-Tahan et. al., 2021).

**Proposition 3.3:** Let  $(\mathcal{N}, +, \cdot)$  be a *NR* and  $\mathcal{A}, \mathcal{B}$  be two intuitionistic multi-fuzzy sub-*NR*s of  $\mathcal{N}$ , then  $\mathcal{A} \cap \mathcal{B}$  is also an intuitionistic multi-fuzzy sub *NR* of  $\mathcal{N}$ .

Decision making under incomplete data: Intuitionistic multi fuzzy ideals of near-ring approach

**Proof.** Let  $\mathcal{A}, \mathcal{B}$  be two intuitionistic multi-fuzzy sub-NRs of  $\mathcal{N}$ , then  $\forall s, t \in \mathcal{N}$ , we are to show conditions of definition 3.1.

$$\begin{aligned} \mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}(s - t) &= \mathcal{CM}_{\mathcal{A}}(s - t) \wedge \mathcal{CM}_{\mathcal{B}}(s - t) \\ &\geq \{\mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{A}}(t)\} \wedge \{\mathcal{CM}_{\mathcal{B}}(s) \wedge \mathcal{CM}_{\mathcal{B}}(t)\} \\ &= \{\mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{B}}(s)\} \wedge \{\mathcal{CM}_{\mathcal{A}}(t) \wedge \mathcal{CM}_{\mathcal{B}}(t)\} \\ &= \mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}(s) \wedge \mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}(t) \Rightarrow \mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}(s - t) \\ &\geq \mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}(s) \wedge \mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}(t) \end{aligned}$$

$$\begin{aligned} \text{And, } \mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}(s - t) &= \mathcal{CN}_{\mathcal{A}}(s - t) \vee \mathcal{CN}_{\mathcal{B}}(s - t) \leq \{\mathcal{CN}_{\mathcal{A}}(s) \vee \mathcal{CN}_{\mathcal{A}}(t)\} \vee \\ &\{\mathcal{CN}_{\mathcal{B}}(s) \vee \mathcal{CN}_{\mathcal{B}}(t)\} = \{\mathcal{CN}_{\mathcal{A}}(s) \vee \mathcal{CN}_{\mathcal{B}}(s)\} \vee \{\mathcal{CN}_{\mathcal{A}}(t) \vee \mathcal{CN}_{\mathcal{B}}(t)\} = \mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}(s) \vee \\ &\mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}(t) \Rightarrow \mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}(s - t) \leq \mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}(s) \vee \mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}(t) \end{aligned}$$

$$\begin{aligned} \text{Also, } \mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}(s \cdot t) &= \mathcal{CM}_{\mathcal{A}}(s \cdot t) \wedge \mathcal{CM}_{\mathcal{B}}(s \cdot t) \geq \{\mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{A}}(t)\} \wedge \\ &\{\mathcal{CM}_{\mathcal{B}}(s) \wedge \mathcal{CM}_{\mathcal{B}}(t)\} = \{\mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{B}}(s)\} \wedge \{\mathcal{CM}_{\mathcal{A}}(t) \wedge \mathcal{CM}_{\mathcal{B}}(t)\} = \mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}(s) \wedge \\ &\mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}(t) \Rightarrow \mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}(s \cdot t) \geq \mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}(s) \wedge \mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}(t) \end{aligned}$$

$$\begin{aligned} \text{And, } \mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}(s \cdot t) &= \mathcal{CN}_{\mathcal{A}}(s \cdot t) \vee \mathcal{CN}_{\mathcal{B}}(s \cdot t) \leq \{\mathcal{CN}_{\mathcal{A}}(s) \vee \mathcal{CN}_{\mathcal{A}}(t)\} \vee \{\mathcal{CN}_{\mathcal{B}}(s) \vee \\ &\mathcal{CN}_{\mathcal{B}}(t)\} = \{\mathcal{CN}_{\mathcal{A}}(s) \vee \mathcal{CN}_{\mathcal{B}}(s)\} \vee \{\mathcal{CN}_{\mathcal{A}}(t) \vee \mathcal{CN}_{\mathcal{B}}(t)\} = \mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}(s) \vee \mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}(t) \Rightarrow \\ &\mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}(s \cdot t) \leq \mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}(s) \vee \mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}(t) \end{aligned}$$

**Corollary 1.** Let  $(\mathcal{N}, +, \cdot)$  be a NR, and  $\mathcal{A}_i$  be an intuitionistic multi-fuzzy sub-NRs of  $\mathcal{N}$  for  $i = 1, 2, 3, \dots, n$ , then  $\cap_{i=1}^n \mathcal{A}_i$  is also an intuitionistic multi-fuzzy sub-NR of  $\mathcal{N}$ .

**Proposition 3.4:** (Al-Tahan et. al., 2021 & Shinoj et. al., 2015) Let  $(\mathcal{N}, +, \cdot)$  be a NR, and  $\mathcal{A}$  and  $\mathcal{B}$  be intuitionistic multi-fuzzy ideals of  $\mathcal{N}$ . Then,  $\mathcal{A} \cap \mathcal{B}$  is also an intuitionistic multi-fuzzy ideal of  $\mathcal{N}$ .

**Proof.** Let  $\mathcal{A}, \mathcal{B}$  be two intuitionistic multi-fuzzy ideals of  $\mathcal{N}$ , then  $\forall s, t \in \mathcal{N}$ , we are to show conditions of definition 3.2. The first two have been done in proposition 3.3.

$$\Rightarrow \mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}(s + t - s) = \mathcal{CM}_{\mathcal{A}}(s + t - s) \wedge \mathcal{CM}_{\mathcal{B}}(s + t - s) \geq \mathcal{CM}_{\mathcal{A}}(t) \wedge \mathcal{CM}_{\mathcal{B}}(t)$$

Since  $\mathcal{A}$  and  $\mathcal{B}$  are IMFIs  $\Rightarrow \mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}(s - t) \geq \mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}(t)$

$$\begin{aligned} \text{And, } \mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}(s + t - s) &= \mathcal{CN}_{\mathcal{A}}(s + t - s) \vee \mathcal{CN}_{\mathcal{B}}(s + t - s) \leq \mathcal{CN}_{\mathcal{A}}(t) \vee \mathcal{CN}_{\mathcal{B}}(t) \\ &\Rightarrow \mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}(s - t) \leq \mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}(t) \end{aligned}$$

$$\begin{aligned} \text{Moreover, } \mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}(s \cdot t) &= \mathcal{CM}_{\mathcal{A}}(s \cdot t) \wedge \mathcal{CM}_{\mathcal{B}}(s \cdot t) \geq \mathcal{CM}_{\mathcal{A}}(t) \wedge \mathcal{CM}_{\mathcal{B}}(t) \\ &\Rightarrow \mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}(s \cdot t) \geq \mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}(t) \end{aligned}$$

$$\begin{aligned} \text{Also, } \mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}(s \cdot t) &= \mathcal{CN}_{\mathcal{A}}(s \cdot t) \vee \mathcal{CN}_{\mathcal{B}}(s \cdot t) \leq \mathcal{CN}_{\mathcal{A}}(t) \vee \mathcal{CN}_{\mathcal{B}}(t) \\ &\Rightarrow \mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}(s \cdot t) \leq \mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}(t) \end{aligned}$$

$$\begin{aligned} \text{Let } a \in \mathcal{U} \text{ then } \mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}((s + a)t - st) &= \mathcal{CM}_{\mathcal{A}}((s + a)t - st) \wedge \mathcal{CM}_{\mathcal{B}}((s + a)t - \\ &st) \geq \mathcal{CM}_{\mathcal{A}}(a) \wedge \mathcal{CM}_{\mathcal{B}}(a) \end{aligned}$$

$$\text{Because } \mathcal{A} \text{ and } \mathcal{B} \text{ are IMFIs } \Rightarrow \mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}((s + a)t - st) \geq \mathcal{CM}_{\mathcal{A} \cap \mathcal{B}}(a)$$

$$\begin{aligned} \text{And, } \mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}((s + a)t - st) &= \mathcal{CN}_{\mathcal{A}}((s + a)t - st) \vee \mathcal{CN}_{\mathcal{B}}((s + a)t - st) \leq \mathcal{CN}_{\mathcal{A}}(a) \vee \\ &\mathcal{CN}_{\mathcal{B}}(a) \Rightarrow \mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}((s + a)t - st) \leq \mathcal{CN}_{\mathcal{A} \cap \mathcal{B}}(a) \quad \blacksquare \end{aligned}$$

**Corollary 2.** Let  $(\mathcal{N}, +, \cdot)$  be a NR, and  $\mathcal{A}_i$  be the intuitionistic multi-fuzzy ideal of NRs of  $\mathcal{N}$  for  $i = 1, 2, 3, \dots, n$  then  $\cap_{i=1}^n \mathcal{A}_i$  is also an intuitionistic multi-fuzzy ideal of  $\mathcal{N}$ .

**Example 9.** Let  $(Z_3, +, \cdot)$  be a NR and  $\mathcal{A}, \mathcal{B}$  be two intuitionistic multi-fuzzy ideals of  $Z_3$  given by;

$$\begin{aligned} \mathcal{A} = \{ < 0, (0.9, 0.9, 0.5, 0.5), (0.1, 0.1, 0.5, 0.5) >, < 1, (0.7, 0.3, 0.1, 0.1), (0.3, 0.7, 0.9, 0.9) \\ >, < 2, (0.7, 0.3, 0.1, 0.1), (0.3, 0.7, 0.9, 0.9) > \} \end{aligned}$$

And

$$\mathcal{B} = \{ \langle 0, (0.7, 0.7, 0.7), (0.3, 0.3, 0.3) \rangle, \langle 1, (0.5, 0.5, 0.2), (0.5, 0.5, 0.8) \rangle, \langle 2, (0.5, 0.5, 0.2), (0.5, 0.5, 0.8) \rangle \}$$

$$\mathcal{A} \cap \mathcal{B} = \{ \langle 0, (0.7, 0.7, 0.5), (0.3, 0.3, 0.5) \rangle, \langle 1, (0.5, 0.3, 0.1), (0.5, 0.7, 0.9) \rangle, \langle 2, (0.5, 0.3, 0.1), (0.5, 0.7, 0.9) \rangle \}$$

It also satisfies the conditions of definition 3.2 and forms an intuitionistic multi-fuzzy ideal of  $\mathbb{Z}_3$ .

**Remark 6.** Let  $(\mathcal{N}, +, \cdot)$  be a NR and  $\mathcal{A}, \mathcal{B}$  be two intuitionistic multi-fuzzy ideals of  $\mathcal{N}$ . Then,  $\mathcal{A} \cup \mathcal{B}$  may or may not be an intuitionistic multi-fuzzy ideal of  $\mathcal{N}$ .

**Example 10.** Let  $(\mathbb{Z}, +, \cdot)$  be a NR and  $\mathcal{A}, \mathcal{B}$  be two intuitionistic multi-fuzzy ideals of  $\mathbb{Z}$  given by;

$$\mathcal{CM}_{\mathcal{A}}(s) = \begin{cases} (0.6, 0.6) & \text{if } s \text{ is multiple of } 5 \\ (0.3, 0.2) & \text{otherwise} \end{cases}$$

$$\mathcal{CN}_{\mathcal{A}}(s) = \begin{cases} (0.4, 0.4) & \text{if } s \text{ is multiple of } 5 \\ (0.7, 0.8) & \text{otherwise} \end{cases}$$

And,

$$\mathcal{CM}_{\mathcal{B}}(s) = \begin{cases} (0.5, 0.4) & \text{if } 2/s \\ (0.4, 0.3) & \text{otherwise} \end{cases}$$

$$\mathcal{CN}_{\mathcal{B}}(s) = \begin{cases} (0.5, 0.6) & \text{if } 2/s \\ (0.6, 0.7) & \text{otherwise} \end{cases}$$

Then,

$$\mathcal{CM}_{\mathcal{A} \cup \mathcal{B}}(s) = \begin{cases} (0.6, 0.6) & \text{if } s \text{ is multiple of } 5 \text{ and } 2/s \\ (0.5, 0.4) & \text{if } 2/s \\ (0.4, 0.3) & \text{otherwise} \end{cases}$$

$$\mathcal{CN}_{\mathcal{A} \cup \mathcal{B}}(s) = \begin{cases} (0.4, 0.4) & \text{if } s \text{ is multiple of } 5 \text{ and } 2/s \\ (0.5, 0.6) & \text{if } 2/s \\ (0.6, 0.7) & \text{otherwise} \end{cases}$$

So,  $\mathcal{CM}_{\mathcal{A} \cup \mathcal{B}}(6 - 5) \not\geq \mathcal{CM}_{\mathcal{A} \cup \mathcal{B}}(6) \wedge \mathcal{CM}_{\mathcal{A} \cup \mathcal{B}}(5)$  M Vble number *ot* of definition 3.2 and form intuitionistic fuzzy multi ideal of  $\mathcal{CN}_{\mathcal{A} \cup \mathcal{B}}(6 - 5) \not\leq \mathcal{CM}_{\mathcal{A} \cup \mathcal{B}}(6) \vee \mathcal{CM}_{\mathcal{A} \cup \mathcal{B}}(5)$ .

It does not satisfy the conditions of definition 3.2 and does not form the intuitionistic multi-fuzzy ideal of  $\mathbb{Z}$ .

**Proposition 3.5:** Let  $(\mathcal{N}, +, \cdot)$  be a NR, and  $\mathcal{A}$  be an intuitionistic multi-fuzzy ideal of  $\mathcal{N}$  (Shinoj et. al., 2015), then;

$$\mathcal{CM}_{\mathcal{A}}(s) \leq \mathcal{CM}_{\mathcal{A}}(0) \text{ and } \mathcal{CN}_{\mathcal{A}}(s) \geq \mathcal{CN}_{\mathcal{A}}(0) \quad \forall s \in \mathcal{N}$$

$$\mathcal{CM}_{\mathcal{A}}(-s) = \mathcal{CM}_{\mathcal{A}}(s) \text{ and } \mathcal{CN}_{\mathcal{A}}(-s) = \mathcal{CN}_{\mathcal{A}}(s) \quad \forall s \in \mathcal{N}$$

$$\mathcal{CM}_{\mathcal{A}}(s + t - s) = \mathcal{CM}_{\mathcal{A}}(t) \text{ and } \mathcal{CN}_{\mathcal{A}}(s + t - s) = \mathcal{CN}_{\mathcal{A}}(t) \quad \forall s, t \in \mathcal{N}$$

**Proof 1).** Let  $s \in \mathcal{N}$ , then we have

$$\mathcal{CM}_{\mathcal{A}}(0) = \mathcal{CM}_{\mathcal{A}}(s - s) \geq \mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{A}}(s) \geq \mathcal{CM}_{\mathcal{A}}(s)$$

$$\text{Also, } \mathcal{CN}_{\mathcal{A}}(0) = \mathcal{CN}_{\mathcal{A}}(s - s) \leq \mathcal{CN}_{\mathcal{A}}(s) \vee \mathcal{CN}_{\mathcal{A}}(s) \leq \mathcal{CN}_{\mathcal{A}}(s)$$

**Proof 2).** Let  $s \in \mathcal{N}$ , then we have

$$\mathcal{CM}_{\mathcal{A}}(-s) = \mathcal{CM}_{\mathcal{A}}(0 - s) \geq \mathcal{CM}_{\mathcal{A}}(0) \wedge \mathcal{CM}_{\mathcal{A}}(s) \geq \mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{A}}(s) = \mathcal{CM}_{\mathcal{A}}(s)$$

$$\text{Also, } \mathcal{CN}_{\mathcal{A}}(-s) = \mathcal{CN}_{\mathcal{A}}(0 - s) \leq \mathcal{CN}_{\mathcal{A}}(0) \vee \mathcal{CN}_{\mathcal{A}}(s) \leq \mathcal{CN}_{\mathcal{A}}(s) \vee \mathcal{CN}_{\mathcal{A}}(s) =$$

$$\mathcal{CN}_{\mathcal{A}}(s)$$



Decision making under incomplete data: Intuitionistic multi fuzzy ideals of near-ring approach

**Proof 3).** Is straight forward

**Proposition 3.6:** Let  $(\mathcal{N}, +, \cdot)$  be a *NR*, and  $\mathcal{A}$  be an intuitionistic multi-fuzzy ideal of  $\mathcal{N}$  if;

$$\mathcal{CM}_{\mathcal{A}}(s - t) = \mathcal{CM}_{\mathcal{A}}(0) \Rightarrow \mathcal{CM}_{\mathcal{A}}(s) = \mathcal{CM}_{\mathcal{A}}(t) \text{ and}$$

$$\mathcal{CN}_{\mathcal{A}}(s - t) = \mathcal{CN}_{\mathcal{A}}(0) \Rightarrow \mathcal{CN}_{\mathcal{A}}(s) = \mathcal{CN}_{\mathcal{A}}(t)$$

**Proof.** Let  $\mathcal{A}$  be an intuitionistic multi-fuzzy ideal of  $\mathcal{N}$  and  $\mathcal{CM}_{\mathcal{A}}(s - t) = \mathcal{CM}_{\mathcal{A}}(0)$   
 $\forall s, t \in \mathcal{N}$  then,

$$\mathcal{CM}_{\mathcal{A}}(s) = \mathcal{CM}_{\mathcal{A}}(s - t + t) \geq \mathcal{CM}_{\mathcal{A}}(s - t) \wedge \mathcal{CM}_{\mathcal{A}}(t) \geq \mathcal{CM}_{\mathcal{A}}(0) \wedge \mathcal{CM}_{\mathcal{A}}(t)$$

Using proposition 3.5

$$, \mathcal{CM}_{\mathcal{A}}(s) = \mathcal{CM}_{\mathcal{A}}(t.)$$

Also, suppose that  $\mathcal{CN}_{\mathcal{A}}(s - t) = \mathcal{CN}_{\mathcal{A}}(0) \forall s, t \in \mathcal{N}$ . Then,

$$\mathcal{CN}_{\mathcal{A}}(s) = \mathcal{CN}_{\mathcal{A}}(s - t + t) \leq \mathcal{CN}_{\mathcal{A}}(s - t) \vee \mathcal{CN}_{\mathcal{A}}(t) \leq \mathcal{CN}_{\mathcal{A}}(0) \vee \mathcal{CN}_{\mathcal{A}}(t)$$

$$\Rightarrow \mathcal{CN}_{\mathcal{A}}(s) = \mathcal{CN}_{\mathcal{A}}(t)$$

**Definition 3.7:** Let  $(\mathcal{N}, +, \cdot)$  be a *NR*, and  $\mathcal{A}$  be an intuitionistic multi-fuzzy set of  $\mathcal{N}$ , then  $\mathcal{A}^*$  is defined as  $\mathcal{A}^* = \{s \in \mathcal{N} \text{ such that } \mathcal{CM}_{\mathcal{A}}(s) = \mathcal{CM}_{\mathcal{A}}(0) \text{ and } \mathcal{CN}_{\mathcal{A}}(s) = \mathcal{CN}_{\mathcal{A}}(0)\}$ .

**Example 11.** Let  $(\mathbb{Z}, +, \cdot)$  be a *NR* and  $\mathcal{A}, \mathcal{B}$  be two intuitionistic multi-fuzzy ideals of  $\mathbb{Z}$  defined in example 10

. Then,  $\mathcal{A}^* = 5\mathbb{Z}$

**Example 12.** Let  $(\mathbb{Z}_3, +, \cdot)$  be a *NR* and  $\mathcal{B}$  be an intuitionistic multi-fuzzy ideal of  $\mathbb{Z}_3$  defined in example 9. Then,  $\mathcal{B}^* = \emptyset$

**Lemma 3.8:** Let  $(\mathcal{N}, +, \cdot)$  be a *NR*, and  $\mathcal{A}$  be an intuitionistic multi-fuzzy ideal of  $\mathcal{N}$  then;

$$\mathcal{CM}_{\mathcal{A}}(s) \leq \mathcal{CM}_{\mathcal{A}}(0) \text{ and } \mathcal{CN}_{\mathcal{A}}(s) \geq \mathcal{CN}_{\mathcal{A}}(0) \forall s \in \mathcal{N}$$

**Proof.** The proof is done in proposition 3.5.

**Proposition 3.9:** Let  $(\mathcal{N}, +, \cdot)$  be a *NR* and  $\mathcal{A}$  be an intuitionistic multi-fuzzy sub-*NR* of  $\mathcal{N}$ , then  $\mathcal{A}^*$  is also sub-*NR* of  $\mathcal{N}$  (Shinoj et. al. , 2015) .

**Proof.** Let  $s, t \in \mathcal{A}^*$  then,

$$\mathcal{CM}_{\mathcal{A}}(s) = \mathcal{CM}_{\mathcal{A}}(0) \text{ and } \mathcal{CN}_{\mathcal{A}}(s) = \mathcal{CN}_{\mathcal{A}}(0)$$

$$\mathcal{CM}_{\mathcal{A}}(t) = \mathcal{CM}_{\mathcal{A}}(0) \text{ and } \mathcal{CN}_{\mathcal{A}}(t) = \mathcal{CN}_{\mathcal{A}}(0)$$

$$\mathcal{CM}_{\mathcal{A}}(s - t) \geq \mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{A}}(t) = \mathcal{CM}_{\mathcal{A}}(0) \text{ and that } \mathcal{CM}_{\mathcal{A}}(s) \leq \mathcal{CM}_{\mathcal{A}}(0) \forall s \in \mathcal{N}$$

$$\Rightarrow \mathcal{CM}_{\mathcal{A}}(s - t) = \mathcal{CM}_{\mathcal{A}}(0)$$

$$\text{Also, } \mathcal{CN}_{\mathcal{A}}(s - t) \leq \mathcal{CN}_{\mathcal{A}}(s) \vee \mathcal{CN}_{\mathcal{A}}(t) = \mathcal{CN}_{\mathcal{A}}(0) \text{ and that } \mathcal{CN}_{\mathcal{A}}(s) \geq \mathcal{CN}_{\mathcal{A}}(0) \forall s \in \mathcal{N}$$

$$\Rightarrow \mathcal{CN}_{\mathcal{A}}(s - t) = \mathcal{CN}_{\mathcal{A}}(0)$$

hence  $s - t \in \mathcal{A}^*$

$$\text{Moreover, } \mathcal{CM}_{\mathcal{A}}(s \cdot t) \geq \mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{A}}(t) = \mathcal{CM}_{\mathcal{A}}(0) \text{ and that } \mathcal{CM}_{\mathcal{A}}(s) \leq \mathcal{CM}_{\mathcal{A}}(0) \forall s \in \mathcal{N}$$

$$\Rightarrow \mathcal{CM}_{\mathcal{A}}(s \cdot t) = \mathcal{CM}_{\mathcal{A}}(0)$$

$$\text{Also, } \mathcal{CN}_{\mathcal{A}}(s \cdot t) \leq \mathcal{CN}_{\mathcal{A}}(s) \vee \mathcal{CN}_{\mathcal{A}}(t) = \mathcal{CN}_{\mathcal{A}}(0) \text{ and that } \mathcal{CN}_{\mathcal{A}}(s) \geq \mathcal{CN}_{\mathcal{A}}(0) \forall s \in \mathcal{N}$$

$$\Rightarrow \mathcal{CN}_{\mathcal{A}}(s \cdot t) = \mathcal{CN}_{\mathcal{A}}(0)$$

hence  $s \cdot t \in \mathcal{A}^*$

**Proposition 3.10:** (Al-Tahan et. al. 2021) Let  $(\mathcal{N}, +, \cdot)$  be a *NR*, and  $\mathcal{A}$  be an intuitionistic multi-fuzzy ideal of  $\mathcal{N}$ , then  $\mathcal{A}^*$  is also an ideal of  $\mathcal{N}$ .

**Proof.** Let  $s \in \mathcal{A}^*$  and  $a, b \in \mathcal{N}$  then,

$$\mathcal{CM}_{\mathcal{A}}(s) = \mathcal{CM}_{\mathcal{A}}(0) \text{ and } \mathcal{CN}_{\mathcal{A}}(s) = \mathcal{CN}_{\mathcal{A}}(0)$$

$$\Rightarrow \mathcal{CM}_{\mathcal{A}}(a + s - a) \geq \mathcal{CM}_{\mathcal{A}}(s) = \mathcal{CM}_{\mathcal{A}}(0) \text{ and that } \mathcal{CM}_{\mathcal{A}}(s) \leq \mathcal{CM}_{\mathcal{A}}(0) \forall s \in \mathcal{N}$$

$$\Rightarrow \mathcal{CM}_{\mathcal{A}}(a + s - a) = \mathcal{CM}_{\mathcal{A}}(0)$$

$$\text{Also, } \mathcal{CN}_{\mathcal{A}}(a + s - a) \leq \mathcal{CN}_{\mathcal{A}}(s) = \mathcal{CN}_{\mathcal{A}}(0) \text{ and that } \mathcal{CN}_{\mathcal{A}}(s) \geq \mathcal{CN}_{\mathcal{A}}(0) \forall s \in \mathcal{N}$$

$$\Rightarrow \mathcal{CN}_{\mathcal{A}}(a + s - a) = \mathcal{CN}_{\mathcal{A}}(0)$$

hence  $(a + s - a) \in \mathcal{A}^*$

$$\text{Moreover, } \mathcal{CM}_{\mathcal{A}}(a \cdot s) \geq \mathcal{CM}_{\mathcal{A}}(s) = \mathcal{CM}_{\mathcal{A}}(0) \text{ and that } \mathcal{CM}_{\mathcal{A}}(s) \leq \mathcal{CM}_{\mathcal{A}}(0) \forall s \in \mathcal{N}$$

$$\Rightarrow \mathcal{CM}_{\mathcal{A}}(a \cdot s) = \mathcal{CM}_{\mathcal{A}}(0)$$

$$\text{Also, } \mathcal{CN}_{\mathcal{A}}(a \cdot s) \leq \mathcal{CN}_{\mathcal{A}}(s) = \mathcal{CN}_{\mathcal{A}}(0) \text{ and that } \mathcal{CN}_{\mathcal{A}}(s) \geq \mathcal{CN}_{\mathcal{A}}(0) \forall s \in \mathcal{N}$$

$$\Rightarrow \mathcal{CN}_{\mathcal{A}}(a \cdot s) = \mathcal{CN}_{\mathcal{A}}(0)$$

hence  $a \cdot s \in \mathcal{A}^*$

$$\text{Now, } \mathcal{CM}_{\mathcal{A}}((a + s)b - ab) \geq \mathcal{CM}_{\mathcal{A}}(s) = \mathcal{CM}_{\mathcal{A}}(0) \text{ and that } \mathcal{CM}_{\mathcal{A}}(s) \leq \mathcal{CM}_{\mathcal{A}}(0) \forall s \in \mathcal{N}$$

$$\Rightarrow \mathcal{CM}_{\mathcal{A}}((a + s)b - ab) = \mathcal{CM}_{\mathcal{A}}(0)$$

$$\text{Also } \mathcal{CN}_{\mathcal{A}}((a + s)b - ab) \leq \mathcal{CN}_{\mathcal{A}}(s) = \mathcal{CN}_{\mathcal{A}}(0) \text{ and that } \mathcal{CN}_{\mathcal{A}}(s) \geq \mathcal{CN}_{\mathcal{A}}(0) \forall s \in \mathcal{N}$$

$$\Rightarrow \mathcal{CN}_{\mathcal{A}}((a + s)b - ab) = \mathcal{CN}_{\mathcal{A}}(0)$$

Hence  $((a + s)b - ab) \in \mathcal{A}^*$

**Definition 3.11:** Let  $\mathcal{N}$  be a non-empty set, and  $\mathcal{A}$  be an intuitionistic multi-fuzzy set of  $\mathcal{N}$ ; then the support of  $\mathcal{N}$  can be defined as  $\mathcal{A}_* = \{s \in \mathcal{N} \text{ such that } \mathcal{CM}_{\mathcal{A}}(s) > 0 \text{ and } \mathcal{CN}_{\mathcal{A}}(s) < 1\}$ .

**Example 13.** Let  $(\mathbb{Z}, +, \cdot)$  be a NR under standard addition and multiplication, and  $\mathcal{A}$  be an intuitionistic multi-fuzzy sub-NR over  $\mathbb{Z}$  defined in example 7, then,  $\mathcal{A}_* = 2\mathbb{Z}$

**Proposition 3.12:** Let  $(\mathcal{N}, +, \cdot)$  be a NR, and  $\mathcal{A}$  be an intuitionistic multi-fuzzy sub NR of  $\mathcal{N}$ , then  $\mathcal{A}_*$  is also sub NR of  $\mathcal{N}$ .

**Proof.** Let  $s, t \in \mathcal{A}_*$ , then,  $\mathcal{CM}_{\mathcal{A}}(s) > 0$  and  $\mathcal{CN}_{\mathcal{A}}(s) < 1$  also  $\mathcal{CM}_{\mathcal{A}}(t) > 0$  and  $\mathcal{CN}_{\mathcal{A}}(t) < 1$

$$\Rightarrow \mathcal{CM}_{\mathcal{A}}(s - t) \geq \mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{A}}(t) > 0 \Rightarrow \mathcal{CM}_{\mathcal{A}}(s - t) > 0$$

$$\text{Also, } \mathcal{CN}_{\mathcal{A}}(s - t) \leq \mathcal{CN}_{\mathcal{A}}(s) \vee \mathcal{CN}_{\mathcal{A}}(t) < 1 \Rightarrow \mathcal{CN}_{\mathcal{A}}(s - t) < 1$$

hence  $s - t \in \mathcal{A}_*$

$$\text{Moreover, } \mathcal{CM}_{\mathcal{A}}(s \cdot t) \geq \mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{A}}(t) > 0 \Rightarrow \mathcal{CM}_{\mathcal{A}}(s \cdot t) > 0$$

$$\text{Also, } \mathcal{CN}_{\mathcal{A}}(s \cdot t) \leq \mathcal{CN}_{\mathcal{A}}(s) \vee \mathcal{CN}_{\mathcal{A}}(t) < 1$$

$$\Rightarrow \mathcal{CN}_{\mathcal{A}}(s \cdot t) < 1 \text{ hence } s \cdot t \in \mathcal{A}_*$$

**Proposition 3.13:** Let  $(\mathcal{N}, +, \cdot)$  be a NR, and  $\mathcal{A}$  be an intuitionistic multi-fuzzy ideal of  $\mathcal{N}$ , then  $\mathcal{A}_*$  is also an ideal of  $\mathcal{N}$ .

**Proof.** Let  $s \in \mathcal{A}_*$  and  $a, b \in \mathcal{N}$  then  $\mathcal{CM}_{\mathcal{A}}(s) > 0$  and  $\mathcal{CN}_{\mathcal{A}}(s) < 1$

$$\Rightarrow \mathcal{CM}_{\mathcal{A}}(a + s - a) \geq \mathcal{CM}_{\mathcal{A}}(s) > 0 \Rightarrow \mathcal{CM}_{\mathcal{A}}(a + s - a) > 0$$

$$\text{Also, } \mathcal{CN}_{\mathcal{A}}(a + s - a) \leq \mathcal{CN}_{\mathcal{A}}(s) < 1 \Rightarrow \mathcal{CN}_{\mathcal{A}}(a + s - a) < 1$$

hence  $(a + s - a) \in \mathcal{A}_*$

$$\text{Moreover, } \mathcal{CM}_{\mathcal{A}}(a \cdot s) \geq \mathcal{CM}_{\mathcal{A}}(s) > 0 \Rightarrow \mathcal{CM}_{\mathcal{A}}(a \cdot s) > 0$$

$$\text{Also, } \mathcal{CN}_{\mathcal{A}}(a \cdot s) \leq \mathcal{CN}_{\mathcal{A}}(s) < 1 \Rightarrow \mathcal{CN}_{\mathcal{A}}(a \cdot s) < 1$$

Decision making under incomplete data: Intuitionistic multi fuzzy ideals of near-ring approach  
hence  $a \cdot s \in \mathcal{A}_*$

Now,  $\mathcal{CM}_{\mathcal{A}}((a + s)b - ab) \geq \mathcal{CM}_{\mathcal{A}}(s) > 0 \Rightarrow \mathcal{CM}_{\mathcal{A}}((a + s)b - ab) > 0$

Also  $\mathcal{CN}_{\mathcal{A}}((a + s)b - ab) \leq \mathcal{CN}_{\mathcal{A}}(s) < 1 \Rightarrow \mathcal{CN}_{\mathcal{A}}((a + s)b - ab) < 1$

Hence  $((a + s)b - ab) \in \mathcal{A}_*$

**Definition 3.14:** Let  $\mathcal{U}$  and  $\mathcal{V}$  be two non-empty sets,  $\mathcal{A}$  and  $\mathcal{B}$  be two intuitionistic fuzzy multisets of  $\mathcal{U}$  and  $\mathcal{V}$ , respectively, then  $\mathcal{A} \times \mathcal{B}$  can be defined as,

$$\mathcal{A} \times \mathcal{B}(u, v) = \{ \langle (u, v), \mathcal{CM}_{\mathcal{A}}(u) \wedge \mathcal{CM}_{\mathcal{B}}(v), \mathcal{CN}_{\mathcal{A}}(u) \vee \mathcal{CN}_{\mathcal{B}}(v) \rangle \}$$

**Proposition 3.15:** Let  $\mathcal{U}$  and  $\mathcal{V}$  be two NRs and  $\mathcal{A}, \mathcal{B}$  be intuitionistic multi-fuzzy sub-NR of  $\mathcal{U}, \mathcal{V}$  respectively, then  $\mathcal{A} \times \mathcal{B}$  is also an intuitionistic multi-fuzzy sub-NR of  $\mathcal{U} \times \mathcal{V}$ .

**Proof.** Let  $\mathcal{A}$  and  $\mathcal{B}$  be two intuitionistic multi-fuzzy sub-NRs and

$(u_1, v_1), (u_2, v_2) \in \mathcal{U} \times \mathcal{V}$ . Then we are to show the conditions of Definition 3.1.

(I)

$$\begin{aligned} \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}((u_1 - u_2), (v_1 - v_2)) &= \mathcal{CM}_{\mathcal{A}}(u_1 - u_2) \wedge \mathcal{CM}_{\mathcal{B}}(v_1 - v_2) \\ &\geq \{ \mathcal{CM}_{\mathcal{A}}(u_1) \wedge \mathcal{CM}_{\mathcal{A}}(u_2) \} \wedge \{ \mathcal{CM}_{\mathcal{B}}(v_1) \wedge \mathcal{CM}_{\mathcal{B}}(v_2) \} \\ &= \{ \mathcal{CM}_{\mathcal{A}}(u_1) \wedge \mathcal{CM}_{\mathcal{B}}(v_1) \} \wedge \{ \mathcal{CM}_{\mathcal{A}}(u_2) \wedge \mathcal{CM}_{\mathcal{B}}(v_2) \} \\ &= \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(u_1, v_1) \wedge \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(u_2, v_2) \\ \Rightarrow \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}((u_1 - u_2), (v_1 - v_2)) &\geq \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(u_1, v_1) \wedge \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(u_2, v_2) \end{aligned}$$

And,

$$\begin{aligned} \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}((u_1 - u_2), (v_1 - v_2)) &= \mathcal{CN}_{\mathcal{A}}(u_1 - u_2) \vee \mathcal{CN}_{\mathcal{B}}(v_1 - v_2) \\ &\leq \{ \mathcal{CN}_{\mathcal{A}}(u_1) \vee \mathcal{CN}_{\mathcal{A}}(u_2) \} \vee \{ \mathcal{CN}_{\mathcal{B}}(v_1) \vee \mathcal{CN}_{\mathcal{B}}(v_2) \} \\ &= \{ \mathcal{CN}_{\mathcal{A}}(u_1) \vee \mathcal{CN}_{\mathcal{B}}(v_1) \} \vee \{ \mathcal{CN}_{\mathcal{A}}(u_2) \vee \mathcal{CN}_{\mathcal{B}}(v_2) \} \\ &= \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(u_1, v_1) \vee \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(u_2, v_2) \\ \Rightarrow \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}((u_1 - u_2), (v_1 - v_2)) &\leq \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(u_1, v_1) \vee \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(u_2, v_2) \end{aligned}$$

(II)

$$\begin{aligned} \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}((u_1 u_2), (v_1 v_2)) &= \mathcal{CM}_{\mathcal{A}}(u_1 u_2) \wedge \mathcal{CM}_{\mathcal{B}}(v_1 v_2) \\ &\geq \{ \mathcal{CM}_{\mathcal{A}}(u_1) \wedge \mathcal{CM}_{\mathcal{A}}(u_2) \} \wedge \{ \mathcal{CM}_{\mathcal{B}}(v_1) \wedge \mathcal{CM}_{\mathcal{B}}(v_2) \} \\ &= \{ \mathcal{CM}_{\mathcal{A}}(u_1) \wedge \mathcal{CM}_{\mathcal{B}}(v_1) \} \wedge \{ \mathcal{CM}_{\mathcal{A}}(u_2) \wedge \mathcal{CM}_{\mathcal{B}}(v_2) \} \\ &= \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(u_1, v_1) \wedge \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(u_2, v_2) \\ \Rightarrow \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}((u_1 u_2), (v_1 v_2)) &\geq \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(u_1, v_1) \wedge \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(u_2, v_2) \end{aligned}$$

And,

$$\begin{aligned} \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}((u_1 u_2), (v_1 v_2)) &= \mathcal{CN}_{\mathcal{A}}(u_1 u_2) \vee \mathcal{CN}_{\mathcal{B}}(v_1 v_2) \\ &\leq \{ \mathcal{CN}_{\mathcal{A}}(u_1) \vee \mathcal{CN}_{\mathcal{A}}(u_2) \} \vee \{ \mathcal{CN}_{\mathcal{B}}(v_1) \vee \mathcal{CN}_{\mathcal{B}}(v_2) \} \\ &= \{ \mathcal{CN}_{\mathcal{A}}(u_1) \vee \mathcal{CN}_{\mathcal{B}}(v_1) \} \vee \{ \mathcal{CN}_{\mathcal{A}}(u_2) \vee \mathcal{CN}_{\mathcal{B}}(v_2) \} \\ &= \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(u_1, v_1) \vee \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(u_2, v_2) \\ \Rightarrow \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}((u_1 u_2), (v_1 v_2)) &\leq \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(u_1, v_1) \vee \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(u_2, v_2) \end{aligned}$$

**Corollary 3.** Let  $\mathcal{N}_i$  be NRs and  $\mathcal{A}_i$  be intuitionistic multi-fuzzy sub-NR of  $\mathcal{N}_i$  for  $i = 1, 2, \dots, n$ , then  $\prod_{i=1}^n \mathcal{A}_i$  is an intuitionistic multi-fuzzy sub-NR of  $\prod_{i=1}^n \mathcal{N}_i$  where  $\mathcal{CM}_{\mathcal{A}_1 \times \mathcal{A}_2 \times \dots \times \mathcal{A}_n}(n_1, n_2, \dots, n_n) = \bigwedge_{i=1}^n \mathcal{CM}_{\mathcal{A}_i}(n_i)$  and  $\mathcal{CN}_{\mathcal{A}_1 \times \mathcal{A}_2 \times \dots \times \mathcal{A}_n}(n_1, n_2, \dots, n_n) = \bigvee_{i=1}^n \mathcal{CN}_{\mathcal{A}_i}(n_i) \forall n_i \in \mathcal{N}_i$ .

**Proposition 3.16:** Let  $\mathcal{U}$  and  $\mathcal{V}$  be two NRs, and  $\mathcal{A}$  and  $\mathcal{B}$  be the intuitionistic multi-fuzzy ideal of  $\mathcal{U}$  and  $\mathcal{V}$ , respectively. Then,  $\mathcal{A} \times \mathcal{B}$  is also an intuitionistic multi-fuzzy ideal of  $\mathcal{U} \times \mathcal{V}$  (Al-Tahan et. al., 2021 & Sujatha, 2104).

**Proof.** Let  $\mathcal{A}$  and  $\mathcal{B}$  be two intuitionistic multi-fuzzy sub-NRs and  $(u_1, v_1), (u_2, v_2) \in \mathcal{U} \times \mathcal{V}$ , then we are to show conditions of Definition 3.2. (I) and (II) has been done in proposition 3.15 then;

$$\begin{aligned} \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}((u_1 + u_2 - u_1), (v_1 + v_2 - v_1)) &= \mathcal{CM}_{\mathcal{A}}(u_1 + u_2 - u_1) \wedge \mathcal{CM}_{\mathcal{B}}(v_1 + v_2 - v_1) \\ &\geq \mathcal{CM}_{\mathcal{A}}(u_2) \wedge \mathcal{CM}_{\mathcal{B}}(v_2) = \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(u_2, v_2) \quad \therefore \mathcal{A}, \mathcal{B} \text{ are IMFI's} \\ \Rightarrow \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}((u_1 + u_2 - u_1), (v_1 + v_2 - v_1)) &\geq \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(u_2, v_2) \end{aligned}$$

$$\begin{aligned} \text{and } \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}((u_1 + u_2 - u_1), (v_1 + v_2 - v_1)) & \\ &= \mathcal{CN}_{\mathcal{A}}(u_1 + u_2 - u_1) \vee \mathcal{CN}_{\mathcal{B}}(v_1 + v_2 - v_1) \\ &\leq \mathcal{CN}_{\mathcal{A}}(u_2) \vee \mathcal{CN}_{\mathcal{B}}(v_2) = \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(u_2, v_2) \end{aligned}$$

$$\Rightarrow \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}((u_1 + u_2 - u_1), (v_1 + v_2 - v_1)) \leq \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(u_2, v_2)$$

$$\text{Also, } \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}((u_1 u_2), (v_1 v_2)) = \mathcal{CM}_{\mathcal{A}}(u_1 u_2) \wedge \mathcal{CM}_{\mathcal{B}}(v_1 v_2) \geq \mathcal{CM}_{\mathcal{A}}(u_2) \wedge \mathcal{CM}_{\mathcal{B}}(v_2) = \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(u_2, v_2) \quad \therefore \mathcal{A}, \mathcal{B} \text{ are IMFI's}$$

$$\Rightarrow \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}((u_1 u_2), (v_1 v_2)) \geq \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(u_2, v_2)$$

$$\text{And, } \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}((u_1 u_2), (v_1 v_2)) = \mathcal{CN}_{\mathcal{A}}(u_1 u_2) \vee \mathcal{CN}_{\mathcal{B}}(v_1 v_2) \leq \mathcal{CN}_{\mathcal{A}}(u_2) \vee \mathcal{CN}_{\mathcal{B}}(v_2) = \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(u_2, v_2)$$

$$\Rightarrow \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}((u_1 u_2), (v_1 v_2)) \leq \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(u_2, v_2)$$

$$\Rightarrow \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}((u_1 u_2), (v_1 v_2)) \leq \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(u_2, v_2)$$

Let  $n_1, n_2 \in \mathcal{N}$  then,

$$\begin{aligned} \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}((u_1 + n_1)u_2 - u_1 u_2, (v_1 + n_2)v_2 - v_1 v_2)) & \\ &= \mathcal{CM}_{\mathcal{A}}((u_1 + n_1)u_2 - u_1 u_2) \wedge \mathcal{CM}_{\mathcal{B}}((v_1 + n_2)v_2 - v_1 v_2) \\ &\geq \mathcal{CM}_{\mathcal{A}}(n_1) \wedge \mathcal{CM}_{\mathcal{B}}(n_2) = \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(n_1, n_2) \end{aligned}$$

$$\Rightarrow \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}((u_1 + n_1)u_2 - u_1 u_2, (v_1 + n_2)v_2 - v_1 v_2)) \geq \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(n_1, n_2)$$

$$\text{And } \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}((u_1 + n_1)u_2 - u_1 u_2, (v_1 + n_2)v_2 - v_1 v_2)) = \mathcal{CN}_{\mathcal{A}}((u_1 + n_1)u_2 - u_1 u_2) \vee \mathcal{CN}_{\mathcal{B}}((v_1 + n_2)v_2 - v_1 v_2) \leq \mathcal{CN}_{\mathcal{A}}(n_1) \vee \mathcal{CN}_{\mathcal{B}}(n_2) = \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(n_1, n_2)$$

$$\Rightarrow \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}((u_1 + n_1)u_2 - u_1 u_2, (v_1 + n_2)v_2 - v_1 v_2)) \leq \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(n_1, n_2)$$

**Corollary 4.** Let  $\mathcal{N}_i$  be NRs and  $\mathcal{A}_i$  be intuitionistic multi-fuzzy ideals of  $\mathcal{N}_i$  for  $i = 1, 2, \dots, n$ , then  $\prod_{i=1}^n \mathcal{A}_i$  is an intuitionistic multi-fuzzy ideal of  $\prod_{i=1}^n \mathcal{N}_i$  where  $\mathcal{CM}_{\mathcal{A}_1 \times \mathcal{A}_2 \times \dots \times \mathcal{A}_n}(n_1, n_2, \dots, n_n) = \bigwedge_{i=1}^n \mathcal{CM}_{\mathcal{A}_i}(n_i)$

and  $\mathcal{CN}_{\mathcal{A}_1 \times \mathcal{A}_2 \times \dots \times \mathcal{A}_n}(n_1, n_2, \dots, n_n) = \bigvee_{i=1}^n \mathcal{CN}_{\mathcal{A}_i}(n_i) \quad \forall n_i \in \mathcal{N}_i$ .

**Example 14.** Let  $(\mathbb{Z}, +, \cdot)$  be a NR and  $\mathcal{A}, \mathcal{B}$  be two intuitionistic multi-fuzzy ideals of  $\mathbb{Z}$  defined in example 10. Then  $\mathcal{A} \times \mathcal{B}$  is an intuitionistic multi-fuzzy ideal of  $\mathbb{Z} \times \mathbb{Z}$  with count functions given by;

$$\mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(s, t) = \begin{cases} (0.5, 0.4) & \text{if } s \text{ is multiple of 5 and } 2/t \\ (0.4, 0.3) & \text{if } s \text{ is multiple of 5} \\ (0.3, 0.2) & \text{otherwise} \end{cases}$$

$$\mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(s, t) = \begin{cases} (0.5, 0.6) & \text{if } s \text{ is multiple of 5 and } 2/t \\ (0.6, 0.7) & \text{if } s \text{ is multiple of 5} \\ (0.7, 0.8) & \text{otherwise} \end{cases}$$

**Definition 3.17:** Let  $\mathcal{N}$  be a non-empty set and  $\mathcal{A}$  be an intuitionistic multi-fuzzy set of  $\mathcal{N}$  and  $\theta = \{\theta_1, \theta_2, \theta_3, \dots, \theta_n\}, \delta = \{\delta_1, \delta_2, \dots, \delta_n\}$  where  $\theta, \delta \in [0, 1]$  with  $\theta + \delta \leq 1$  then level subset of an intuitionistic multi fuzzy set  $\mathcal{A}$  can be defined as;  
 $(\mathcal{A})_{\theta, \delta} = \{s \in \mathcal{N} \mid \mathcal{CM}_{\mathcal{A}}(s) \geq \theta \text{ and } \mathcal{CN}_{\mathcal{A}}(s) \leq \delta\}$

Decision making under incomplete data: Intuitionistic multi fuzzy ideals of near-ring approach

**Proposition 3.18:** Let  $\mathcal{N}$  be a non-empty set, and  $\mathcal{A}$  and  $\mathcal{B}$  be intuitionistic multi-fuzzy sets of  $\mathcal{N}$ , then the following results hold;

$(\mathcal{A})_{\theta,\delta} \subseteq (\mathcal{A})_{\gamma,\emptyset}$ , if  $\theta \geq \gamma$  and  $\delta \leq \emptyset$

$$(\mathcal{A})_{1-\delta,\delta} \subseteq (\mathcal{A})_{\theta,\delta} \subseteq (\mathcal{A})_{\theta,1-\theta}$$

$$\mathcal{A} \subseteq \mathcal{B} \Rightarrow (\mathcal{A})_{\theta,\delta} \subseteq (\mathcal{B})_{\theta,\delta}$$

$$(\mathcal{A} \cap \mathcal{B})_{\theta,\delta} = (\mathcal{A})_{\theta,\delta} \cap (\mathcal{B})_{\theta,\delta}$$

$$(\cap \mathcal{A}_i)_{\theta,\delta} = \cap (\mathcal{A}_i)_{\theta,\delta}$$

$$(\mathcal{A})_{0,1} = \mathcal{N}$$

$$(\mathcal{A} \cup \mathcal{B})_{\theta,\delta} \supseteq (\mathcal{A})_{\theta,\delta} \cup (\mathcal{B})_{\theta,\delta}$$

**Remark 7.** Equality hold for  $\forall$  if  $\theta + \delta = 1$ .

**Theorem 3.19:** (Sujatha, 2104) Let  $(\mathcal{N}, +, \cdot)$  be a NR and  $\mathcal{A}$  be an intuitionistic multi-fuzzy sub-NR of  $\mathcal{N}$  then  $(\mathcal{A})_{\theta,\delta}$  is sub-NR of  $\mathcal{N} \quad \forall \theta, \delta \in [0,1]$  with  $\theta + \delta \leq 1$  and  $\mathcal{CM}_{\mathcal{A}}(0) \geq \theta, \mathcal{CN}_{\mathcal{A}}(0) \leq \delta$ .

**Proof.** Given that  $\mathcal{CM}_{\mathcal{A}}(0) \geq \theta, \mathcal{CN}_{\mathcal{A}}(0) \leq \delta$  and

let  $\mathcal{A}$  is an IMF sub - NR of  $\mathcal{N}$  and  $s, t \in (\mathcal{A})_{\theta,\delta}$

$$\Rightarrow \mathcal{CM}_{\mathcal{A}}(s) \geq \theta, \mathcal{CN}_{\mathcal{A}}(s) \leq \delta \text{ and } \mathcal{CM}_{\mathcal{A}}(t) \geq \theta, \mathcal{CN}_{\mathcal{A}}(t) \leq \delta$$

$$\Rightarrow \mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{A}}(t) \geq \theta \text{ and } \mathcal{CN}_{\mathcal{A}}(s) \vee \mathcal{CN}_{\mathcal{A}}(t) \leq \delta$$

$$\Rightarrow \mathcal{CM}_{\mathcal{A}}(s - t) \geq \mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{A}}(t) \geq \theta \text{ and } \mathcal{CN}_{\mathcal{A}}(s - t) \leq \mathcal{CN}_{\mathcal{A}}(s) \vee \mathcal{CN}_{\mathcal{A}}(t) \leq \delta$$

Hence  $s - t \in (\mathcal{A})_{\theta,\delta}$

$$\text{Also, } \mathcal{CM}_{\mathcal{A}}(s \cdot t) \geq \mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{A}}(t) \geq \theta \text{ and } \mathcal{CN}_{\mathcal{A}}(s \cdot t) \leq \mathcal{CN}_{\mathcal{A}}(s) \vee \mathcal{CN}_{\mathcal{A}}(t) \leq \delta$$

Hence  $s \cdot t \in (\mathcal{A})_{\theta,\delta}$

**Theorem 3.20:** Let  $(\mathcal{N}, +, \cdot)$  be a NR and  $\mathcal{A}$  be an intuitionistic multi-fuzzy ideal of  $\mathcal{N}$  then  $(\mathcal{A})_{\theta,\delta}$  is an ideal of  $\mathcal{N} \quad \forall \theta, \delta \in [0,1]$  with  $\theta + \delta \leq 1$  and  $\mathcal{CM}_{\mathcal{A}}(0) \geq \theta, \mathcal{CN}_{\mathcal{A}}(0) \leq \delta$ .

**Proof.** Given that  $\mathcal{CM}_{\mathcal{A}}(0) \geq \theta, \mathcal{CN}_{\mathcal{A}}(0) \leq \delta$  and

let  $\mathcal{A}$  be an IMF I of  $\mathcal{N}$  and  $s, t \in (\mathcal{A})_{\theta,\delta}$  and  $a, b \in \mathcal{N}$

$\Rightarrow \mathcal{CM}_{\mathcal{A}}(s) \geq \theta, \mathcal{CN}_{\mathcal{A}}(s) \leq \delta$  and  $\mathcal{CM}_{\mathcal{A}}(t) \geq \theta, \mathcal{CN}_{\mathcal{A}}(t) \leq \delta$ . Then we are to show the conditions of Definition 3.2.

$$\Rightarrow \mathcal{CM}_{\mathcal{A}}(a + t - a) \geq \mathcal{CM}_{\mathcal{A}}(t) \geq \theta \text{ and } \mathcal{CN}_{\mathcal{A}}(a + t - a) \leq \mathcal{CN}_{\mathcal{A}}(t) \leq \delta$$

Hence  $a + t - a \in (\mathcal{A})_{\theta,\delta}$

$$\text{Also, } \mathcal{CM}_{\mathcal{A}}(a \cdot t) \geq \mathcal{CM}_{\mathcal{A}}(t) \geq \theta \text{ and } \mathcal{CN}_{\mathcal{A}}(a \cdot t) \leq \mathcal{CN}_{\mathcal{A}}(t) \leq \delta$$

Hence  $a \cdot t \in (\mathcal{A})_{\theta,\delta}$

$$\text{Now, } \mathcal{CM}_{\mathcal{A}}((a + t)b - ab) \geq \mathcal{CM}_{\mathcal{A}}(t) \geq \theta \text{ and } \mathcal{CN}_{\mathcal{A}}((a + t)b - ab) \leq \mathcal{CN}_{\mathcal{A}}(t) \leq \delta$$

Hence  $((a + t)b - ab) \in (\mathcal{A})_{\theta,\delta}$

**Definition 3.21:** If  $\mathcal{A} \times \mathcal{B}$  is the Cartesian product of two intuitionistic fuzzy multisets and  $\theta = \{\theta_1, \theta_2, \theta_3, \dots, \theta_n\}, \delta = \{\delta_1, \delta_2, \dots, \delta_n\}$  where  $\theta, \delta \in [0,1]$  with  $\theta + \delta \leq 1$  then level subset of  $\mathcal{A} \times \mathcal{B}$  can be defined as;

$$(\mathcal{A} \times \mathcal{B})_{\theta,\delta} = \{(s, t) \mid \mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{B}}(t) \geq \theta \text{ and } \mathcal{CN}_{\mathcal{A}}(s) \vee \mathcal{CN}_{\mathcal{B}}(t) \leq \delta\}$$

**Proposition 3.22:** Let  $\mathcal{N}$  be a non-empty set and  $\mathcal{A}, \mathcal{B}$  be two intuitionistic fuzzy multisets of  $\mathcal{N}$ , then  $(\mathcal{A} \times \mathcal{B})_{\theta,\delta} = (\mathcal{A})_{\theta,\delta} \times (\mathcal{B})_{\theta,\delta}$

**Proof.** Let  $(s, t) \in (\mathcal{A})_{\theta,\delta} \times (\mathcal{B})_{\theta,\delta} \Leftrightarrow s \in (\mathcal{A})_{\theta,\delta}$  and  $t \in (\mathcal{B})_{\theta,\delta} \Leftrightarrow \mathcal{CM}_{\mathcal{A}}(s) \geq$

$$\theta \text{ and } \mathcal{CN}_{\mathcal{A}}(s) \leq \delta \text{ also, } \mathcal{CM}_{\mathcal{B}}(t) \geq \theta \text{ and } \mathcal{CN}_{\mathcal{B}}(t) \leq \delta \Leftrightarrow \{\mathcal{CM}_{\mathcal{A}}(s) \wedge \mathcal{CM}_{\mathcal{B}}(t)\} \geq$$

$$\theta \text{ and } \{\mathcal{CN}_{\mathcal{A}}(s) \vee \mathcal{CN}_{\mathcal{B}}(t)\} \leq \delta \Leftrightarrow (s, t) \in (\mathcal{A} \times \mathcal{B})_{\theta,\delta}$$

Hence  $(\mathcal{A} \times \mathcal{B})_{\theta, \delta} = (\mathcal{A})_{\theta, \delta} \times (\mathcal{B})_{\theta, \delta}$

**Theorem 3.23:** Let  $(\mathcal{N}, +, \cdot)$  be a NR and  $\mathcal{A}, \mathcal{B}$  be two intuitionistic multi-fuzzy sub-NR of  $\mathcal{N}$ , and  $\mathcal{A} \times \mathcal{B}$  is an intuitionistic multi-fuzzy sub-NR of  $\mathcal{N} \times \mathcal{N}$  then  $(\mathcal{A} \times \mathcal{B})_{\theta, \delta}$  is sub-NR of  $\mathcal{N} \times \mathcal{N}$ .

**Proof.** Let  $\mathcal{A} \times \mathcal{B}$  be an IMF sub NR of  $\mathcal{N} \times \mathcal{N}$  and  $(s_1, t_1), (s_2, t_2) \in (\mathcal{A} \times \mathcal{B})_{\theta, \delta}$  then;

$$\Rightarrow \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(s_1, t_1) = \mathcal{CM}_{\mathcal{A}}(s_1) \wedge \mathcal{CM}_{\mathcal{B}}(t_1) \geq \theta \text{ and } \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(s_1, t_1) \\ = \mathcal{CN}_{\mathcal{A}}(s_1) \vee \mathcal{CN}_{\mathcal{B}}(t_1) \geq \delta$$

$$\text{similarly, } \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(s_2, t_2) = \mathcal{CM}_{\mathcal{A}}(s_2) \wedge \mathcal{CM}_{\mathcal{B}}(t_2) \geq \theta \text{ and } \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(s_2, t_2) \\ = \mathcal{CN}_{\mathcal{A}}(s_2) \vee \mathcal{CN}_{\mathcal{B}}(t_2) \geq \delta$$

$$\Rightarrow \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}\{(s_1, t_1) - (s_2, t_2)\} \geq \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(s_1, t_1) \wedge \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(s_2, t_2) \\ \geq \theta \text{ and } \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}\{(s_1, t_1) - (s_2, t_2)\} \leq \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(s_1, t_1) \vee \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(s_2, t_2) \\ \leq \delta$$

$$\Rightarrow \{(s_1, t_1) - (s_2, t_2)\} \in (\mathcal{A} \times \mathcal{B})_{\theta, \delta}$$

$$\text{Also, } \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}\{(s_1, t_1) \cdot (s_2, t_2)\} \geq \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(s_1, t_1) \wedge \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(s_2, t_2) \\ \geq \theta \text{ and } \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}\{(s_1, t_1) \cdot (s_2, t_2)\} \leq \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(s_1, t_1) \vee \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(s_2, t_2) \\ \leq \delta$$

$$\Rightarrow \{(s_1, t_1) \cdot (s_2, t_2)\} \in (\mathcal{A} \times \mathcal{B})_{\theta, \delta}$$

**Theorem 3.24:** Let  $(\mathcal{N}, +, \cdot)$  be a NR and  $\mathcal{A}, \mathcal{B}$  be two intuitionistic multi-fuzzy ideals of  $\mathcal{N}$  and  $\mathcal{A} \times \mathcal{B}$  is an intuitionistic multi-fuzzy ideal of  $\mathcal{N} \times \mathcal{N}$ . Then,  $(\mathcal{A} \times \mathcal{B})_{\theta, \delta}$  is an ideal of  $\mathcal{N} \times \mathcal{N}$  (Sujatha, 2014).

**Proof.** Let  $\mathcal{A} \times \mathcal{B}$  be an IMFI of  $\mathcal{N} \times \mathcal{N}$  and  $(s_1, t_1) \in (\mathcal{A} \times$

$\mathcal{B})_{\theta, \delta}$   $(a_1, b_1), (a_2, b_2) \in \mathcal{N} \times \mathcal{N}$  then;  $\mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(s_1, t_1) \geq \theta, \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(s_1, t_1) \leq \delta$

$$\Rightarrow \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}\{(a_1, b_1) + (s_1, t_1) - (a_1, b_1)\} \geq \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(s_1, t_1) \geq \theta$$

$$\text{And, } \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}\{(a_1, b_1) + (s_1, t_1) - (a_1, b_1)\} \leq \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(s_1, t_1) \leq \delta$$

$$\Rightarrow \{(a_1, b_1) + (s_1, t_1) - (a_1, b_1)\} \in (\mathcal{A} \times \mathcal{B})_{\theta, \delta}$$

$$\text{Also, } \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}\{(a_1, b_1) \cdot (s_1, t_1)\} \geq \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(s_1, t_1) \geq \theta \text{ and } \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}\{(a_1, b_1) \cdot (s_1, t_1)\} \leq \\ \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(s_1, t_1) \leq \delta$$

$$\Rightarrow \{(a_1, b_1) \cdot (s_1, t_1)\} \in (\mathcal{A} \times \mathcal{B})_{\theta, \delta}$$

$$\text{Also, } \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}\{((a_1, b_1) + (s_1, t_1))(a_2, b_2) - (a_1, b_1)(a_2, b_2)\} \geq \mathcal{CM}_{\mathcal{A} \times \mathcal{B}}(s_1, t_1) \geq \theta$$

$$\text{And } \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}\{((a_1, b_1) + (s_1, t_1))(a_2, b_2) - (a_1, b_1)(a_2, b_2)\} \leq \mathcal{CN}_{\mathcal{A} \times \mathcal{B}}(s_1, t_1) \leq \delta$$

$$\Rightarrow \{((a_1, b_1) + (s_1, t_1))(a_2, b_2) - (a_1, b_1)(a_2, b_2)\} \in (\mathcal{A} \times \mathcal{B})_{\theta, \delta}$$

**Lemma 3.25:** Let  $(\mathcal{N}, +, \cdot)$  be a NR, and  $\mathcal{A}$  be an intuitionistic multi-fuzzy ideal of  $\mathcal{N}$  then;  $\mathcal{CM}_{\mathcal{A}}(a_1 + a_2 + \dots + a_n) \geq \bigwedge_{1 \leq i \leq n} \mathcal{CM}_{\mathcal{A}}(a_i)$  and  $\mathcal{CN}_{\mathcal{A}}(a_1 + a_2 + \dots + a_n) \leq \bigvee_{1 \leq i \leq n} \mathcal{CN}_{\mathcal{A}}(a_i) \quad \forall a_i \in \mathcal{N}$

**Theorem 3.26:** Let  $(\mathcal{N}, +, \cdot)$  be a NR under standard addition and multiplication is defined as  $a \cdot b = b \quad \forall a, b \in \mathcal{N}$ , and  $\mathcal{A}$  be an intuitionistic multi-fuzzy ideal of  $\mathcal{N}$ .

Then,  $\mathcal{A}'$  is an intuitionistic multi-fuzzy ideal of (matrix near ring)  $\mathcal{M}_n(\mathcal{N})$  where

$$\mathcal{CM}_{\mathcal{A}'}$$
 and  $\mathcal{CN}_{\mathcal{A}'}$  can be defined as;  $\mathcal{CM}_{\mathcal{A}'}([a_{ij}]) =$

$$\bigwedge_{1 \leq i, j \leq n} \mathcal{CM}_{\mathcal{A}}(a_{ij}) \text{ and } \mathcal{CN}_{\mathcal{A}'}([a_{ij}]) = \bigvee_{1 \leq i, j \leq n} \mathcal{CN}_{\mathcal{A}}(a_{ij}) \quad \forall a_{ij}, b_{ij} \in \mathcal{M}_n(\mathcal{N}).$$

**Proof.** Let  $[a_{ij}], [b_{ij}] \in \mathcal{M}_n(\mathcal{N})$  then we are to show conditions of Definition 3.2.

$$\mathcal{CM}_{\mathcal{A}'}([a_{ij}] - [b_{ij}]) = \bigwedge_{1 \leq i, j \leq n} \mathcal{CM}_{\mathcal{A}}(a_{ij} - b_{ij}) \geq \bigwedge_{1 \leq i, j \leq n} \{\mathcal{CM}_{\mathcal{A}}(a_{ij}) \wedge$$

$$\mathcal{CM}_{\mathcal{A}}(b_{ij})\} = \{\bigwedge_{1 \leq i, j \leq n} \mathcal{CM}_{\mathcal{A}}(a_{ij})\} \wedge \{\bigwedge_{1 \leq i, j \leq n} \mathcal{CM}_{\mathcal{A}}(b_{ij})\} = \mathcal{CM}_{\mathcal{A}'}([a_{ij}]) \wedge$$

$$\mathcal{CM}_{\mathcal{A}'}([b_{ij}])$$

Decision making under incomplete data: Intuitionistic multi fuzzy ideals of near-ring approach

$$\begin{aligned}
& \text{And } \mathcal{CN}_{\mathcal{A}'}([a_{ij}] - [b_{ij}]) = \bigvee_{1 \leq i, j \leq n} \mathcal{CN}_{\mathcal{A}}(a_{ij} - b_{ij}) \leq \\
& \bigvee_{1 \leq i, j \leq n} \{\mathcal{CN}_{\mathcal{A}}(a_{ij}) \vee \mathcal{CN}_{\mathcal{A}}(b_{ij})\} = \{\bigvee_{1 \leq i, j \leq n} \mathcal{CN}_{\mathcal{A}}(a_{ij})\} \vee \{\bigvee_{1 \leq i, j \leq n} \mathcal{CN}_{\mathcal{A}}(b_{ij})\} = \\
& \mathcal{CN}_{\mathcal{A}'}([a_{ij}]) \vee \mathcal{CN}_{\mathcal{A}'}([b_{ij}]) \\
& \text{(II) and (IV) of definition 3.2; Since } [a_{ij}] \cdot [b_{ij}] = [c_{ij}] \text{ where } c_{ij} = b_{1j} + b_{2j} + \dots + \\
& b_{nj} \Rightarrow \mathcal{CM}_{\mathcal{A}'}([a_{ij}] \cdot [b_{ij}]) = \mathcal{CM}_{\mathcal{A}'}([c_{ij}]) = \bigwedge_{1 \leq i, j \leq n} \mathcal{CM}_{\mathcal{A}}(c_{ij}) = \\
& \bigwedge_{1 \leq i, j \leq n} \mathcal{CM}_{\mathcal{A}}(c_{1j}) \text{ but } \mathcal{CM}_{\mathcal{A}}(c_{1j}) \geq \mathcal{CM}_{\mathcal{A}}(b_{1j}) \wedge \mathcal{CM}_{\mathcal{A}}(b_{2j}) \wedge \dots \wedge \mathcal{CM}_{\mathcal{A}}(b_{nj}) \Rightarrow \\
& \mathcal{CM}_{\mathcal{A}'}([a_{ij}] \cdot [b_{ij}]) \geq \mathcal{CM}_{\mathcal{A}}(b_{1j}) \wedge \mathcal{CM}_{\mathcal{A}}(b_{2j}) \wedge \dots \wedge \mathcal{CM}_{\mathcal{A}}(b_{nj}) = \\
& \bigwedge_{1 \leq i, j \leq n} \mathcal{CM}_{\mathcal{A}}(b_{ij}) = \mathcal{CM}_{\mathcal{A}'}([b_{ij}]) \\
& \text{And } \mathcal{CN}_{\mathcal{A}'}([a_{ij}] \cdot [b_{ij}]) = \mathcal{CN}_{\mathcal{A}'}([c_{ij}]) = \bigvee_{1 \leq i, j \leq n} \mathcal{CN}_{\mathcal{A}}(c_{ij}) = \\
& \bigvee_{1 \leq i, j \leq n} \mathcal{CN}_{\mathcal{A}}(c_{1j}) \text{ but } \mathcal{CN}_{\mathcal{A}}(c_{1j}) \leq \mathcal{CN}_{\mathcal{A}}(b_{1j}) \vee \mathcal{CN}_{\mathcal{A}}(b_{2j}) \vee \dots \vee \mathcal{CN}_{\mathcal{A}}(b_{nj}) \Rightarrow \\
& \mathcal{CN}_{\mathcal{A}'}([a_{ij}] \cdot [b_{ij}]) \leq \mathcal{CN}_{\mathcal{A}}(b_{1j}) \vee \mathcal{CN}_{\mathcal{A}}(b_{2j}) \vee \dots \vee \mathcal{CN}_{\mathcal{A}}(b_{nj}) = \bigvee_{1 \leq i, j \leq n} \mathcal{CN}_{\mathcal{A}}(b_{ij}) = \\
& \mathcal{CN}_{\mathcal{A}'}([b_{ij}]) \\
& \text{Also, } \mathcal{CM}_{\mathcal{A}'}([a_{ij}] + [b_{ij}] - [a_{ij}]) = \bigwedge_{1 \leq i, j \leq n} \mathcal{CM}_{\mathcal{A}}(a_{ij} + b_{ij} - a_{ij}) \geq \\
& \bigwedge_{1 \leq i, j \leq n} \mathcal{CM}_{\mathcal{A}}(b_{ij}) = \mathcal{CM}_{\mathcal{A}'}([b_{ij}]) \text{ and } \mathcal{CN}_{\mathcal{A}'}([a_{ij}] + [b_{ij}] - [a_{ij}]) = \\
& \bigvee_{1 \leq i, j \leq n} \mathcal{CN}_{\mathcal{A}}(a_{ij} + b_{ij} - a_{ij}) \leq \bigvee_{1 \leq i, j \leq n} \mathcal{CN}_{\mathcal{A}}(b_{ij}) = \mathcal{CN}_{\mathcal{A}'}([b_{ij}]) \\
& \text{Let } [c_{ij}] \in \mathcal{M}_n(\mathcal{N}) \text{ and } \{([a_{ij}] + [b_{ij}])[c_{ij}] - [a_{ij}][c_{ij}]\} = [d_{ij}] \text{ where } d_{ij} = 0 \vee 1 \leq \\
& i, j \leq n \text{ then } \mathcal{CM}_{\mathcal{A}'}([d_{ij}]) = \bigwedge_{1 \leq i, j \leq n} \mathcal{CM}_{\mathcal{A}}(d_{ij}) = \mathcal{CM}_{\mathcal{A}}(0) \geq \mathcal{CM}_{\mathcal{A}}(b_{ij}) = \\
& \mathcal{CM}_{\mathcal{A}'}([b_{ij}]) \\
& \text{And } \mathcal{CN}_{\mathcal{A}'}([d_{ij}]) = \bigvee_{1 \leq i, j \leq n} \mathcal{CN}_{\mathcal{A}}(d_{ij}) = \mathcal{CN}_{\mathcal{A}}(0) \leq \mathcal{CN}_{\mathcal{A}}(b_{ij}) = \mathcal{CN}_{\mathcal{A}'}([b_{ij}])
\end{aligned}$$

**Example 15.** Let  $(\mathbb{Z}, +, \cdot)$  be a NR and  $\mathcal{B}$  be an intuitionistic multi fuzzy ideal of  $\mathbb{Z}$  defined in example 10 then

$$\begin{aligned}
\mathcal{CM} \left( \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \right) &= \begin{cases} (0.5, 0.4) & \text{if } [a_{ij}] \text{ is divisible by 2} \\ (0.4, 0.3) & \text{otherwise} \end{cases} \\
\mathcal{CN} \left( \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \right) &= \begin{cases} (0.5, 0.6) & \text{if } [a_{ij}] \text{ is divisible by 2} \\ (0.6, 0.7) & \text{otherwise} \end{cases}
\end{aligned}$$

is an intuitionistic multi fuzzy ideal of  $\mathcal{M}_2(\mathbb{Z})$ .

**Theorem 3.27:** According to Al-Tahan et. al. (2021), let  $(\mathcal{N}, +, \cdot)$  be a NR under standard addition and multiplication is defined as  $a \cdot b = b \forall a, b \in \mathcal{N}$  and  $\mathcal{A}$  be an intuitionistic multi-fuzzy ideal of  $\mathcal{N}$ . Then,  $\mathcal{A}'$  is an intuitionistic multi-fuzzy ideal of  $P_n(\mathcal{N})$ . where  $\mathcal{CM}_{\mathcal{A}'}$  and  $\mathcal{CN}_{\mathcal{A}'}$  can be defined as;  $\mathcal{CM}_{\mathcal{A}'}(\hbar_0 + \hbar_1s + \hbar_2s^2 + \dots + \hbar_ns^n) = \bigwedge_{0 \leq i \leq n} \mathcal{CM}_{\mathcal{A}}(\hbar_i)$  and  $\mathcal{CN}_{\mathcal{A}'}(\hbar_0 + \hbar_1s + \hbar_2s^2 + \dots + \hbar_ns^n) = \bigvee_{0 \leq i \leq n} \mathcal{CN}_{\mathcal{A}'}(\hbar_i) \forall \hbar(s) \in P_n(\mathcal{N})$ .

**Proof.** Let  $\hbar(s), g(s) \in P_n(\mathcal{N})$  where  $\hbar(s) = \hbar_0 + \hbar_1s + \hbar_2s^2 + \dots + \hbar_ns^n, g(s) = g_0 + g_1s + g_2s^2 + \dots + g_ns^n$  then we are to show conditions of Definition 3.2.

$$\text{(I) } \mathcal{CM}_{\mathcal{A}'}(\hbar(s) - g(s)) = \bigwedge_{0 \leq i \leq n} \mathcal{CM}_{\mathcal{A}}(\hbar_i - g_i) \geq \bigwedge_{0 \leq i \leq n} \{\mathcal{CM}_{\mathcal{A}}(\hbar_i) \wedge \mathcal{CM}_{\mathcal{A}}(g_i)\} = \\
\{\bigwedge_{0 \leq i \leq n} \mathcal{CM}_{\mathcal{A}}(\hbar_i)\} \wedge \{\bigwedge_{0 \leq i \leq n} \mathcal{CM}_{\mathcal{A}}(g_i)\} = \mathcal{CM}_{\mathcal{A}'}(\hbar(s)) - \mathcal{CM}_{\mathcal{A}'}(g(s))$$

$$\text{And } \mathcal{CN}_{\mathcal{A}'}(\hbar(s) - g(s)) = \bigvee_{0 \leq i \leq n} \mathcal{CN}_{\mathcal{A}}(\hbar_i - g_i) \leq \bigvee_{0 \leq i \leq n} \{\mathcal{CN}_{\mathcal{A}}(\hbar_i) \vee \mathcal{CN}_{\mathcal{A}}(g_i)\} = \\
\{\bigvee_{0 \leq i \leq n} \mathcal{CN}_{\mathcal{A}}(\hbar_i)\} \vee \{\bigvee_{0 \leq i \leq n} \mathcal{CN}_{\mathcal{A}}(g_i)\} = \mathcal{CN}_{\mathcal{A}'}(\hbar(s)) - \mathcal{CN}_{\mathcal{A}'}(g(s))$$

$$\text{(II) and (IV) } \hbar(s) \cdot g(s) = g(s) \Rightarrow \mathcal{CM}_{\mathcal{A}'}(\hbar(s) \cdot g(s)) = \mathcal{CM}_{\mathcal{A}'}(g(s)) \geq \mathcal{CM}_{\mathcal{A}'}(g(s))$$

$$\text{And } \mathcal{CN}_{\mathcal{A}'}(\hbar(s) \cdot g(s)) = \mathcal{CN}_{\mathcal{A}'}(g(s)) \leq \mathcal{CN}_{\mathcal{A}'}(g(s))$$

$$(III), \mathcal{CM}_{\mathcal{A}'}(\hbar(s) + g(s) - \hbar(s)) = \bigwedge_{0 \leq i \leq n} \mathcal{CM}_{\mathcal{A}}(\hbar_i + g_i - \hbar_i) \geq \bigwedge_{0 \leq i \leq n} \mathcal{CM}_{\mathcal{A}}(g_i) = \mathcal{CM}_{\mathcal{A}'}(g(s)) \text{ and } \mathcal{CN}_{\mathcal{A}'}(\hbar(s) + g(s) - \hbar(s)) = \bigvee_{0 \leq i \leq n} \mathcal{CN}_{\mathcal{A}}(\hbar_i + g_i - \hbar_i) \leq \bigvee_{0 \leq i \leq n} \mathcal{CN}_{\mathcal{A}}(g_i) = \mathcal{CN}_{\mathcal{A}'}(g(s))$$

$$(V), \text{ Let } u(s) \in P_n(\mathcal{N}) \text{ where } u(s) = u_0 + u_1s + u_2s^2 + \dots + u_ns^n \text{ and } ((\hbar(s) + g(s))u(s) - \hbar(s)g(s)) = 0 \Rightarrow \mathcal{CM}_{\mathcal{A}}(0) \geq \mathcal{CM}_{\mathcal{A}}(u_i) \geq \bigwedge_{0 \leq i \leq n} \mathcal{CM}_{\mathcal{A}}(u_i) = \mathcal{CM}_{\mathcal{A}'}(u(s))$$

$$\text{And, } \mathcal{CN}_{\mathcal{A}}(0) \leq \mathcal{CN}_{\mathcal{A}}(u_i) \leq \bigvee_{0 \leq i \leq n} \mathcal{CN}_{\mathcal{A}}(u_i) = \mathcal{CN}_{\mathcal{A}'}(u(s))$$

**Example 16.** Let  $(\mathbb{Z}, +, \cdot)$  be a NR and  $\mathcal{B}$  be an intuitionistic multi-fuzzy ideal of  $\mathbb{Z}$  defined in example 10 then;

$$\mathcal{CM}(\hbar_0 + \hbar_1s + \hbar_2s^2 + \hbar_3s^3) = \begin{cases} (0.5, 0.4) & \text{if } \hbar_0, \hbar_1, \hbar_2, \hbar_3 \text{ are divisible by 2} \\ (0.4, 0.3) & \text{otherwise} \end{cases}$$

$$\mathcal{CN}(\hbar_0 + \hbar_1s + \hbar_2s^2 + \hbar_3s^3) = \begin{cases} (0.5, 0.6) & \text{if } \hbar_0, \hbar_1, \hbar_2, \hbar_3 \text{ are divisible by 2} \\ (0.6, 0.7) & \text{otherwise} \end{cases}$$

It is an intuitionistic multi-fuzzy ideal of  $P_3(\mathcal{N})$ .

**Proposition 3.28.** Let  $(\mathcal{N}, +, \cdot)$  be a NR, and  $\mathcal{A}$  be an intuitionistic multi-fuzzy ideal of  $\mathcal{N}$ , then  $\mathcal{A}'$  is an intuitionistic multi-fuzzy ideal of  $P_n(\mathcal{N})$  where

$$\mathcal{CM}_{\mathcal{A}'} \text{ and } \mathcal{CN}_{\mathcal{A}'} \text{ can be defined as; } \mathcal{CM}_{\mathcal{A}'}(\hbar_0 + \hbar_1s + \hbar_2s^2 + \dots + \hbar_ns^n) = \mathcal{CM}_{\mathcal{A}}(\hbar_0) \text{ and } \mathcal{CN}_{\mathcal{A}'}\left(\hbar_0 + \hbar_1s + \hbar_2s^2 + \dots + \hbar_n\right)_{s^n} = \mathcal{CN}_{\mathcal{A}}(\hbar_0) \forall \hbar(s) \in P_n(\mathcal{N}).$$

**Proof.** Same as theorem 3.27.

## 5. Conclusion

This study constructed the concept of intuitionistic multi-fuzzy near-rings and intuitionistic multi-fuzzy ideals. It explored and illustrated some properties related to intuitionistic multi-fuzzy near-rings and intuitionistic multi-fuzzy ideals. Moreover, it investigated the support, level subsets and Cartesian product of intuitionistic multi-fuzzy near-rings and ideals. It established results associated with all these new constructions. This work contributes to fuzzy set theory, widely used in multi-criteria decision-making and pattern recognition problems. In the future, one may extend these notions to AI-related decision-making and pattern recognition research. Alternatively, extending to inter-valued fuzzy systems or applying the intuitionistic multi-fuzzy idea to vector spaces and modules is possible.

**Author Contributions:** B.N. and H.S. developed the theoretical formalism, performed the analytic calculations, and performed the numerical simulations; K.N. contributed to the final version of the manuscript; M.M. and K.S. contributed to the design and implementation of the research, to the analysis of the results and to the writing of the manuscript; L.R.Y.M. wrote and revised the manuscript. All authors discussed the results and commented on the manuscript.

**Data Availability Statement:** No data were used to support this study.

**Conflicts of Interest:** The authors declare that there are no conflicts of interest regarding the publication of this article.



## References

- Abbas, S., Hussain, Z., Hussain, S., Sharif, R., & Hussain, S. (2021). Intuitionistic fuzzy entropy and its applications to multicriteria decision making with IF-TODIM. *Journal of mechanics of continua and mathematical sciences*, 16(7), 99–119. <https://doi.org/10.26782/jmcs.2021.07.00009>.
- Abdullah, L. (2013). Fuzzy multi criteria decision making and its applications: A brief review of category. *Procedia-Social and Behavioral Sciences*, 97, 131–136.
- Abou-Zaid, S. (1991). On fuzzy subnear-rings and ideals. *Fuzzy Sets and Systems*, 44(1), 139–146.
- Al Tahan, M., Hoskova-Mayerova, S., & Davvaz, B. (2021). An approach to fuzzy multi-ideals of near rings. *Journal of Intelligent & Fuzzy Systems*, Preprint, 1–11. <https://doi.org/10.3233/JIFS-202914>.
- Al-Husban, A. (2021). Multi-fuzzy hypergroups. *Italian Journal of Pure and Applied Mathematics*, 46, 382–390.
- Ashraf, A., Ullah, K., Hussain, A., & Bari, M. (2022). Interval-Valued Picture Fuzzy Maclaurin Symmetric Mean Operator with application in Multiple Attribute Decision-Making. *Reports in Mechanical Engineering*, 3(1), 301–317.
- Asif, A., Aydi, H., Arshad, M., Rehman, A., & Tariq, U. (2020). Picture Fuzzy Ideals of Near-Rings. *Journal of Mathematics*, 2020. <https://doi.org/10.1155/2020/8857459>.
- Atanassov, K. (1986). Intuitionistic fuzzy sets. *Fuzzy sets and systems* 20 (1), 87-96.
- Broumi, S., Ajay, D., Chellamani, P., Malayalan, L., Talea, M., Bakali, A., Schweizer, P., & Jafari, S. (2022). Interval Valued Pentapartitioned Neutrosophic Graphs with an Application to MCDM. *Operational Research in Engineering Sciences: Theory and Applications*, 5(3), 68–91.
- Broumi, S., Sundareswaran, R., Shanmugapriya, M., Nordo, G., Talea, M., Bakali, A., & Smarandache, F. (2022). Interval-valued fermatean neutrosophic graphs. *Decision Making: Applications in Management and Engineering*, 5(2), 176–200.
- Dresher, M., & Ore, O. (1938). Theory of Multigroups. *American Journal of Mathematics*, 60(3), 705–733. <https://doi.org/10.2307/2371606>
- Fathi, M., & Salleh, A. R. (2009). Intuitionistic fuzzy groups. *Asian Journal of Algebra*, 2(1), 1–10.
- Gulzar, M., Alghazzawi, D., Mateen, M. H., & Kausar, N. (2020). A certain class of t-intuitionistic fuzzy subgroups. *IEEE Access*, 8, 163260–163268. <https://doi.org/10.1109/ACCESS.2020.3020366>.
- Gulzar, M., Mateen, M. H., Alghazzawi, D., & Kausar, N. (2020). A novel application of complex intuitionistic fuzzy sets in group theory. *IEEE Access*, 8, 196075–196085. <https://doi.org/10.1109/ACCESS.2020.3034626>.
- Hoskova-Mayerova, S., & Al Tahan, M. (2021). Anti-Fuzzy Multi-Ideals of Near Ring. *Mathematics*, 9(5), 494. <https://doi.org/10.3390/math8020244>.
- Hur, K., Jang, S.-Y., & Kang, H.-W. (2005). Intuitionistic fuzzy ideals of a ring. *The Pure and Applied Mathematics*, 12(3), 193–209.

Hussain, S., Hussain, S., Rahaman, S., & Abbas, S. (2022). Fuzzy Normed Near-rings and its Ideals. *Journal of Xi'an Shiyou University, Natural Science Edition*, 18(9), 32–36.

Kahraman, C. (2008). Multi-criteria decision-making methods and fuzzy sets. In *Fuzzy multi-criteria decision making*, 1–18. Springer. [https://doi.org/10.1007/978-0-387-76813-7\\_1](https://doi.org/10.1007/978-0-387-76813-7_1).

Kausar, N. (2019). Direct product of finite intuitionistic anti fuzzy normal subrings over non-associative rings. *European Journal of Pure and Applied Mathematics*, 12(2), 622–648.

Kausar, N., Alesemi, M., & Munir, M. (2020). Characterizations of Non-Associative Ordered Semigroups by Their Intuitionistic Fuzzy Bi-Ideals. *Discontinuity, Nonlinearity, and Complexity*, 9(2), 257–275.

Kausar, N., & Waqar, M. A. (2019). Characterizations of non-associative rings by their intuitionistic fuzzy bi-ideals. *European Journal of Pure and Applied Mathematics*, 12(1), 226–250.

Kousar, S., Aslam, F., Kausar, N., & Gaba, Y. U. (2021). Semigroup of finite-state deterministic intuitionistic fuzzy automata with application in fault diagnosis of an aircraft twin-spool turbofan engine. *Journal of Function Spaces*, 2021, 1–10. <https://doi.org/10.1155/2021/1994732>.

Kousar, S., Saleem, T., Kausar, N., Pamucar, D., & Addis, G. M. (2022). Homomorphisms of Lattice-Valued Intuitionistic Fuzzy Subgroup Type-3. *Computational Intelligence and Neuroscience*, 2022, 1–11. <https://doi.org/10.1155/2022/6847138>.

Rahman, S., & Saikia, H. K. (2012). Some aspects of Atanassov's intuitionistic fuzzy submodules. *Int. J. Pure and Appl. Mathematics*, 77(3), 369–383.

Riaz, A., Kousar, S., Kausar, N., Pamucar, D., & Addis, G. (2022a). Codes over Lattice-Valued Intuitionistic Fuzzy Set Type-3 with Application to the Complex DNA Analysis. *Complexity*, 2022, 1–12. <https://doi.org/10.1155/2022/5288187>

Riaz, A., Kousar, S., Kausar, N., Pamucar, D., & Addis, G. M. (2022b). An Analysis of Algebraic Codes over Lattice Valued Intuitionistic Fuzzy Type-3-Submodules. *Computational Intelligence and Neuroscience*, 2022, 1–13. <https://doi.org/10.1155/2022/8148284>

Shah, T., Kausar, N., & Rehman, I. (2012). Intuitionistic fuzzy normal subrings over a non-associative ring. *Analele Universitatii "Ovidius" Constanta-Seria Matematica*, 20(1), 369–386.

Shinoj, T. K., Baby, A., & Sunil, J. J. (2015). On some algebraic structures of fuzzy multisets. *Annals of Fuzzy Mathematics and Informatics*, 9(1), 77–90.

Sujatha, L. (2014). Multi fuzzy subrings and ideals. *Annals of Fuzzy Mathematics and Informatics*, 8(3), 385–392.

Taghieh, A., Mohammadzadeh, A., Zhang, C., Kausar, N., & Castillo, O. (2022). A type-3 fuzzy control for current sharing and voltage balancing in microgrids. *Applied Soft Computing*, 129, 1–13. <https://doi.org/10.1016/j.asoc.2022.109636>.

Decision making under incomplete data: Intuitionistic multi fuzzy ideals of near-ring approach  
Yager, R. R. (1986). On the theory of bags. *International Journal of General System*, 13(1), 23–37.

Zadeh, L. A. (1965). Fuzzy sets. *Information and Control*, 8(3), 338–353.

Zhan, J., & Ma, X. (2005). Intuitionistic fuzzy ideals of near-rings. *Scientiae Mathematicae Japonicae*, 61(2), 219–223.

Zhumadillayeva, A., Orazbayev, B., Santeyeva, S., Dyussekeyev, K., Li, R. Y. M., Crabbe, M. J. C., & Yue, X.-G. (2020). Models for oil refinery waste management using determined and fuzzy conditions. *Information*, 11(6), 299.  
<https://doi.org/10.3390/info11060299>.



© 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).